

MICHIGAN INFANT DEATH STATISTICS
January 1 through December 31, 2017

LANSING, MICHIGAN
NOVEMBER 2018



Rick Snyder, Governor
Nick Lyon, Director

Acknowledgments

DIVISION FOR VITAL RECORDS AND HEALTH STATISTICS

Glenn Copeland, M.B.A. State Registrar

VITAL RECORDS & HEALTH DATA DEVELOPMENT SECTION

Tami Weaver, B.A., Manager

AUTHORS

Glenn Radford, M.A., Statistician

The Division for Vital Records and Health Statistics gratefully acknowledges the contributions of: Kyle Johnson, Kim Essex, Thu Le and Lindsey Myers.



Table Of Contents

Summary	1
---------------	---

Selected Infant Death Statistics, Numbers, Averages & Rates

Table 1	Number of Infant Deaths, Live Births & Infant Death Rates for Michigan Residents, 1970-2017	5
Figure 1	Infant Mortality Rates & Forecasts, 1970-2030	7
Table 2	Number of Infant Deaths, Live Births & Infant Death Rates By Race, 1970-2017	8
Figure 2	Comparison of White & Black Infant Death Rates for Michigan Residents, 1970-2017	10
Table 3	Infant Death Rates by County of Residence, 2017 & 2013 - 2017 Average	11
Table 4	Infant Death Rates by Selected Cities of Residence, 2017 & 2013 - 2017 Average	14
Table 5	Infant Deaths by Race for Michigan, Detroit City & Wayne County, 1990- 2017	15
Table 6	Number & Rate of Infant Deaths by Race, Michigan & United States Residents, 1989 - 2017	19
Table 7-A	Number of Hispanic Deaths by Selected County & Michigan, 2007 - 2017	20
Table 7-B	Number of Middle Easterner Infant Deaths by Selected County & Michigan, 2007 - 2017	21
Table 8	Three-year Average Infant Death Rates by Race & Ancestry, 1999-2017	22
Figure 3	Three-year Average Infant Mortality Rates by Race & Ancestry 1998-2017	23

Perinatal

Table 9	Infant, Hebdomadal, Fetal & Perinatal Death Rates by Specified Race & Ancestry, 2017	24
Table 10	Trends of Infant, Hebdomadal, Fetal & Perinatal Death Rates, 2007-2017	25

Characteristics of Birth and Causes of Death

Table 11	Infant Death Rates by Selected Characteristics of Newborn & of Mother, 2017	26
Table 12	Number of Infant Deaths by Cause of Death, 1980-2017	28
Table 13	Three-year Moving Average Infant Deaths Rates by Cause of Death, 1980-2017	29
Figure 4	Three-year Average Infant Death Rates by Cause, 1980-2017	30
Table 14-A	Dollfus Categories, 2014 - 2017	31
Table 14-B	Dollfus Subcategories, 2014 - 2017	32
Table 15-A	Sudden Unexpected Infant Deaths, 1989 - 2017	33
Table 15-B	Accidental Suffocation & Strangulation, 1989 - 2017	34
Table 16	Sudden Unexpected Infant Deaths by Race, 1989 - 2017	35

Table Of Contents

Three-year Moving Averages by Region

Table 17-A Three-Year Moving Average Infant Death Rates by County, 2006-2017	36
Table 17-B White Three-Year Moving Average Infant Death Rates by County, 2006-2017	39
Table 18-C Black Three-Year Moving Average Infant Death Rates by County, 2006-2017	42
Table 18-A Three-Year Moving Average Infant Death Rates by District, 2006-2017	43
Table 19-B White Three-Year Moving Average Infant Death Rates by District, 2006-2017	45
Table 19-C Black Three-Year Moving Average Infant Death Rates by District, 2006-2017	47
Table 20-A Three-Year Moving Average Infant Death Rates by City, 2006-2017	48
Table 20-B White Three-Year Moving Average Infant Death Rates by City, 2006-2017	49
Table 20-C Black Three-Year Moving Average Infant Death Rates by City, 2006-2017	50

Neonatal & Postneonatal Deaths by Region

Table 21-A Neonatal Deaths by Michigan County & Detroit City , 2007-2017	51
Table 21-B Postneonatal Deaths by Michigan County & Detroit City, 2007-2017	55
Table 22-A Neonatal Deaths by Michigan Health Departments, 2007-2017	59
Table 22-B Postneonatal Deaths by Michigan Health Departments, 2007-2017	61

SUMMARY OF 2017 INFANT DEATH STATISTICS

November 2018 Release

INFANT MORTALITY: For every 1,000 Michigan live births, almost seven infants die before reaching their first birthday. In 2017, 762 infants under the age of one year died, resulting in an infant mortality rate of 6.8 per 1,000 live births. Michigan experienced a significant decline in infant mortality during the early 1990s; but during the 2000s the infant mortality rates remained around 7.9 deaths per 1,000 births. The 2017 rate indicates a drop of the mortality rates to about 7.0 since 2010, from the 7.9 average in the previous decade. The total number of Michigan infants who died between 2010-2017 was 6,200. (See Table 1.)

The perinatal death rate has not significantly changed in years; the rate has been approximately 9 fetal or hebdomadal deaths per 100,000 live births since around 2011. The perinatal death rate was 8.9 in 2016 and 9.3 in 2017. (See Table 10.)

Historically, the infant death rate has declined over time. In 1970, the infant death rate was 20.3 deaths per 1,000 live births; and this rate declined to 10.7 in 1990 and then again to 7.1 deaths per 1,000 live births in 2010. Since 1970, these declines were primarily due to advances in neonatal medicine, artificial lung surfactants, folic acid supplementation, and numerous public health interventions. The gradual decline suggests that the infant mortality rate will drop to about 6.3 by 2020 and perhaps as low as 5.0 by 2030. (See Figure 1.)

Both the white and black infant mortality rates remained about the same in recent years, with a persistent racial disparity in which African American mother's experienced 3 times the risk of an infant death compared to white mothers. In 2005, the white infant death rate was 5.5, and was comparable to the previous ten-year 1996-2005 average of 6.0 deaths per 1,000 white births. The black infant death rate was 17.9 in 2005, and was also comparable to the previous decade average of 17.6 deaths per 1,000 black births. In 2017 the white infant mortality rate was 4.8 per 1,000 live births while the black rate was 14.6 per 1,000 live births. Between 2006-2017 the white infant mortality rate has remained about the same, while the black rate has declined by 25% due to a reduction of infant deaths since 2005. The decline in the Michigan infant mortality rate is largely due to a reduction in black infant deaths. (See Table 2.)

Between 2000-2013, the Native American infant death rate remained on average 11 deaths per 1,000 live births. Between 2015-2017, the rate increased to 14.2 deaths per 1,000 births. The Native American infant death rate is comparable to the African American rate; the racial disparity between Native Americans and whites is about the same as African Americans (2.8 Native American deaths for every white infant death). (See Table 8.)

The Michigan infant mortality rate continues to be higher than the last reported national rate. The 2016 infant death rate for the United States is 5.9. (See Table 6.)

LIVE BIRTHS: In 2017, the number of live births decreased to 111,507, from the 2016 live birth count of 113,374. The 2012-2017 numbers represent historic lows, and Michigan's birth count has gradually decreased since at least 1990. Nationally, there were 3,855,500 births in the U.S. in 2017, a decline of 10.7% from the record number reported for 2007 and a 2.3% decrease from last year.

CAUSES OF DEATH: In 2017, 33.5% of infants died due to conditions related to prematurity and 18.5% died due to birth defects. In addition, 10.8% of infants deaths died due to accidents; 7.1% of all infant deaths were due to accidental suffocation in bed. (See Table 14-A.)

Respiratory Distress Syndrome (RDS) declined from about 15 per 1,000 live births in the early eighties to about 5 per 1,000 by the late nineties. Due to surfactant and other therapies, the number of RDS deaths has continued to decline to less than 2 per 1,000 live births in 2015-2017. In the past few years, the SIDS death rate has remained about 3 per 1,000 live births.

CHARACTERISTICS OF NEWBORN: Certain newborns are at higher risk of dying. In 2017 infants born with very low birth weight (less than 1,500 grams) experienced an infant death rate of 246.0 per 1,000 live births compared to a rate of 2.3 for those infants weighing 2,500 grams or more. Multiple birth infants had an infant mortality rate of 28.4 per 1,000 live births compared to the rate of 6.0 for single birth infants. The excess male infant mortality normally reported in other years was present in 2017, with the infant mortality for male infants at 7.5 per 1,000 male live births and females at 6.1 per 1,000 female live births. (See Table 11.)

CHARACTERISTICS OF MOTHER: Infant death rates were the lowest for mothers aged 30-39 years old and highest for mothers aged under 20 years old. Unmarried mothers had infant mortality rates about twice those of married mothers. Women receiving inadequate prenatal care experienced infant mortality rates three times as high as those women receiving adequate prenatal care. Mothers exposed to secondhand smoking while pregnant had an infant death rate of 9.1 per 1,000 live births compared to a rate of 5.9 for mothers who were not exposed to secondhand smoking during pregnancy. (See Table 11.)

MARCH 2018 UPDATE: In March 2018, the Michigan Division for Vital Records and Health Statistics (DVRHS) reported that 730 Michigan resident infants died in 2016. With the November release, the number of infant deaths in 2016 was updated to 747. Certain mortality records are finalized very late due to forensic enquiries, or for administrative purposes, or for other exceptional reasons. In the case of this 2016 update, the 17 infant deaths were reported late due to exceptional circumstances; this high number of unreported infant deaths is unlikely to occur in 2017.

COMMUNITY LEVEL DATA: Infant mortality data are available for most communities in Michigan. On the Internet, go to Michigan's community health information site at www.michigan.gov/healthstatistics. Data for local health department districts, counties and major cities and townships can be reached by choosing "Vital Statistics" and then "Infant Mortality" button. Those who do not have access to the Internet may request that community level data be mailed to them from the address below.

Questions regarding Infant Death Statistics should be addressed to:

Lindsey Myers
Division for Vital Records and Health Statistics
Michigan Department of Health & Human Services
333 S. Grand Ave.
Lansing, MI 48933

Voice: (517) 335-8715 ♦ FAX: (517) 335-8711 ♦ E-mail:
MyersL@Michigan.gov

Residence Determined by Reported Residence and Geocoded Residence in the Michigan Death Certificate Registry

Using a process called geocoding, the Division for Vital Records & Health Statistics (DVRHS) has assigned geographic coordinates to most death certificates reported between 2000 and 2017. While the primary purpose of geocoding mortality data is to map patterns of death and to conduct spatial epidemiology, such records can also be used to improve the accuracy of location-related data recorded on the certificate. In mortality datasets, the division has found that the MCD residence is miscoded in certain areas about 10-20% of the time. Often, the person reporting the residence confuses the township with an adjacent city or village. Such errors then alter the calculation of local area statistics; small errors in coding lead to large errors in rates calculated for MCDs with populations below 70,000.

Therefore, all mortality and natality statistics between 2000 and 2017 in division reports of infant deaths were updated and recalculated using the geocoded data. **Birth and death counts and rates are different from data reported prior to 2011.** Statistics for years before 2000 use the residence reported during the course of registering a vital event. When rates calculated from these data differ substantially from Michigan or county rates, one should study the events as part of a trend of geocoded statistics, and consider whether the data reflects jumps indicative of a longterm trend of residence-miscoding.

Table 1
Number of Infant Deaths
Live Births and Infant Death Rates
Michigan Residents, 1970 - 2017


Year	Infant Deaths	Live Births	Infant Death Rate
1970	3,492	171,667	20.3 ±0.7
1971	3,157	162,244	19.5 ±0.7
1972	2,801	146,854	19.1 ±0.7
1973	2,561	141,550	18.1 ±0.7
1974	2,387	137,414	17.4 ±0.7
1975	2,205	133,931	16.5 ±0.7
1976	1,978	131,378	15.1 ±0.7
1977	1,945	138,416	14.1 ±0.6
1978	1,931	138,802	13.9 ±0.6
1979	1,921	144,452	13.3 ±0.6
1980	1,851	145,162	12.8 ±0.6
1981	1,851	140,579	13.2 ±0.6
1982	1,672	137,950	12.1 ±0.6
1983	1,573	133,026	11.8 ±0.6
1984	1,595	135,782	11.7 ±0.6
1985	1,575	138,052	11.4 ±0.6
1986	1,565	137,626	11.4 ±0.6
1987	1,538	140,466	10.9 ±0.5
1988	1,542	139,635	11.0 ±0.5
1989	1,645	148,164	11.1 ±0.5
1990	1,638	153,080	10.7 ±0.5
1991	1,554	149,478	10.4 ±0.5
1992	1,460	143,827	10.2 ±0.5
1993	1,319	139,560	9.5 ±0.5
1994	1,184	137,844	8.6 ±0.5
1995	1,110	134,169	8.3 ±0.5

Table 1
Number of Infant Deaths
Live Births and Infant Death Rates
Michigan Residents, 1970 - 2017
(Continued)

Year	Infant Deaths	Live Births	Infant Death Rate
1996	1,072	133,231	8.0 ±0.5
1997	1,085	133,549	8.1 ±0.5
1998	1,091	133,649	8.2 ±0.5
1999	1,071	133,429	8.0 ±0.5
2000	1,112	136,048	8.2 ±0.5
2001	1,066	133,247	8.0 ±0.5
2002	1,054	129,518	8.1 ±0.5
2003	1,112	130,850	8.5 ±0.5
2004	984	129,710	7.6 ±0.5
2005	1,013	127,518	7.9 ±0.5
2006	940	127,537	7.4 ±0.5
2007	997	125,172	8.0 ±0.5
2008	894	121,231	7.4 ±0.5
2009	881	117,309	7.5 ±0.5
2010	817	114,717	7.1 ±0.5
2011	749	114,159	6.6 ±0.5
2012	783	112,708	6.9 ±0.5
2013	799	113,732	7.0 ±0.5
2014	773	114,460	6.8 ±0.5
2015	770	113,211	6.8 ±0.5
2016	747	113,374	6.6 ±0.5
2017	762	111,507	6.8 ±0.5

Note: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Source: 1970 - 2017 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services



Note: Rates are per 1,000 live births. Rates after 2017 are forecasted within an 80% prediction interval indicated by the dotted lines; this means that the future rates fall between the lower and upper bounds of the interval with 80% probability.

Source: 1970- 2017 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 2
Number of Infant Deaths, Live Births and Infant Death Rates By Race
Michigan Residents, 1970 - 2017

Year	White			Black			Other ¹		
	Infant Deaths	Live Births	Infant Death Rate	Infant Deaths	Live Births	Infant Death Rate	Infant Deaths	Live Births	Infant Death Rate
1970	2,661	144,159	18.5 ±0.7	811	26,513	30.6 ±2.1	14	871	16.1 ±8.4
1971	2,293	134,486	17.1 ±0.7	843	26,755	31.5 ±2.1	17	890	19.1 ±9.0
1972	1,957	120,042	16.3 ±0.7	835	25,760	32.4 ±2.2	8	936	8.5 ±5.9
1973	1,749	115,150	15.2 ±0.7	796	25,252	31.5 ±2.2	11	926	11.9 ±7.0
1974	1,673	112,768	14.8 ±0.7	705	22,836	30.9 ±2.2	7	950	7.4 ±5.4
1975	1,574	110,058	14.3 ±0.7	616	22,692	27.1 ±2.1	8	951	8.4 ±5.8
1976	1,383	108,295	12.8 ±0.7	575	21,778	26.4 ±2.1	16	1,100	14.5 ±7.1
1977	1,373	114,511	12.0 ±0.6	559	22,620	24.7 ±2.0	8	1,059	7.6 ±5.2
1978	1,344	114,652	11.7 ±0.6	566	22,423	25.2 ±2.1	13	1,507	8.6 ±4.7
1979	1,352	119,218	11.3 ±0.6	550	23,346	23.6 ±1.9	12	1,622	7.4 ±4.2
1980	1,246	119,698	10.4 ±0.6	582	23,422	24.8 ±2.0	15	1,825	8.2 ±4.1
1981	1,261	116,596	10.8 ±0.6	564	22,002	25.6 ±2.1	18	1,847	9.7 ±4.5
1982	1,110	114,465	9.7 ±0.6	548	21,574	25.4 ±2.1	13	1,797	7.2 ±3.9
1983	1,048	109,892	9.5 ±0.6	506	21,295	23.8 ±2.0	15	1,721	8.7 ±4.4
1984	1,034	111,574	9.3 ±0.6	541	22,322	24.2 ±2.0	15	1,792	8.4 ±4.2
1985	1,053	114,150	9.2 ±0.6	510	21,935	23.3 ±2.0	11	1,820	6.0 ±3.6
1986	999	112,255	8.9 ±0.5	553	23,366	23.7 ±1.9	12	1,927	6.2 ±3.5
1987	953	113,386	8.4 ±0.5	567	24,975	22.7 ±1.8	17	1,992	8.5 ±4.0
1988	952	112,232	8.5 ±0.5	570	25,217	22.6 ±1.8	19	2,093	9.1 ±4.1
1989	946	115,186	8.2 ±0.5	672	29,839	22.5 ±1.7	23	2,525	9.1 ±3.7
1990	930	118,180	7.9 ±0.5	689	31,842	21.6 ±1.6	18	2,623	6.9 ±3.2
1991	857	114,983	7.5 ±0.5	683	31,543	21.7 ±1.6	13	2,522	5.2 ±2.8
1992	781	111,458	7.0 ±0.5	656	29,656	22.1 ±1.7	23	2,223	10.3 ±4.2
1993	770	108,329	7.1 ±0.5	530	28,249	18.8 ±1.6	17	2,418	7.0 ±3.3
1994	670	106,828	6.3 ±0.5	495	27,129	18.2 ±1.6	19	3,137	6.1 ±2.7
1995	655	105,274	6.2 ±0.5	432	24,914	17.3 ±1.6	19	3,124	6.1 ±2.7
1996	627	104,922	6.0 ±0.5	421	24,073	17.5 ±1.7	21	3,307	6.4 ±2.7
1997	640	104,980	6.1 ±0.5	425	24,201	17.6 ±1.7	17	3,628	4.7 ±2.2
1998	660	105,161	6.3 ±0.5	405	24,179	16.8 ±1.6	23	3,833	6.0 ±2.4
1999	616	104,493	5.9 ±0.5	428	23,850	17.9 ±1.7	19	4,210	4.5 ±2.0

Table 2
Number of Infant Deaths, Live Births and Infant Death Rates By Race
Michigan Residents, 1970 - 2017

Year	White			Black			Other ¹		
	Infant Deaths	Live Births	Infant Death Rate	Infant Deaths	Live Births	Infant Death Rate	Infant Deaths	Live Births	Infant Death Rate
2000	637	106,322	6.0 ±0.5	437	24,069	18.2 ±1.7	30	4,525	6.6 ±2.4
2001	635	104,489	6.1 ±0.5	398	23,494	16.9 ±1.7	25	4,752	5.3 ±2.1
2002	611	101,611	6.0 ±0.5	410	22,248	18.4 ±1.8	21	5,056	4.2 ±1.8
2003	680	102,146	6.7 ±0.5	391	22,380	17.5 ±1.7	37	5,508	6.7 ±2.2
2004	527	100,795	5.2 ±0.4	388	22,484	17.3 ±1.7	61	5,719	10.7 ±2.7
2005	549	99,117	5.5 ±0.5	400	22,365	17.9 ±1.7	62	5,461	11.4 ±2.8
2006	534	98,551	5.4 ±0.5	339	22,873	14.8 ±1.6	60	5,636	10.6 ±2.7
2007	549	93,868	5.8 ±0.5	368	22,314	16.5 ±1.7	74	6,932	10.7 ±2.4
2008	496	91,509	5.4 ±0.5	332	22,695	14.6 ±1.6	61	6,691	9.1 ±2.3
2009	470	87,471	5.4 ±0.5	346	22,292	15.5 ±1.6	46	6,980	6.6 ±1.9
2010	462	84,597	5.5 ±0.5	312	22,046	14.2 ±1.6	38	7,425	5.1 ±1.6
2011	411	84,172	4.9 ±0.5	300	21,950	13.7 ±1.5	34	7,529	4.5 ±1.5
2012	455	82,949	5.5 ±0.5	291	21,623	13.5 ±1.5	35	7,696	4.5 ±1.5
2013	473	83,339	5.7 ±0.5	290	22,055	13.1 ±1.5	35	8,081	4.3 ±1.4
2014	452	84,492	5.3 ±0.5	284	21,282	13.3 ±1.5	36	8,574	4.2 ±1.4
2015	423	82,938	5.1 ±0.5	311	21,457	14.5 ±1.6	22	8,583	2.6 ±1.1
2016	401	83,388	4.8 ±0.5	294	21,523	13.7 ±1.6	33	8,182	4.0 ±1.4
2017	392	80,967	4.8 ±0.5	320	21,897	14.6 ±1.6	35	8,222	4.3 ±1.4


Note: Infant deaths by race of infant; live births used in calculating infant death rates are by race of mother. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Infant deaths of unknown race are not included in this table.

¹Between 2004-2008, multi-racial infants were reported as "other."

Source: 1970 - 2017 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Figure 2
Comparison of White and Black Infant Death Rates for Michigan Residents, 1970-2017



Note: Rates are per 1,000 live births. The range indicated by the grey band represents a 95% confidence interval; this means that the true rate lies between the lower and upper bounds of the interval with 95% statistical confidence.

Source: 1970-2017 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 3
Number of Infant Deaths, Live Births and Infant Death Rates by
Michigan and Michigan County of Residence,
2017 and 2013 - 2017 Average

County of Residence	2017			2013 - 2017		
	Infant Deaths	Live Births	Infant Death Rate	Average Infant Deaths	Average Live Births	Average Infant Death Rate
MICHIGAN	762	111,507	6.8 ±0.5	770.2	113,256.8	6.8 ±0.2
Alcona	1	58	*	0.4	59.0	*
Alger	-	63	-	-	62.0	-
Allegan	4	1,348	*	7.4	1,374.0	5.4 ±1.7
Alpena	2	290	*	1.2	271.4	4.4 ±3.5
Antrim	-	161	-	0.6	192.6	*
Arenac	-	113	-	1.2	123.8	9.7 ±7.7
Baraga	1	69	*	1.0	72.6	*
Barry	5	635	*	2.6	633.8	4.1 ±2.2
Bay	7	1,011	6.9 ±5.1	5.4	1,035.4	5.2 ±2.0
Benzie	1	159	*	1.2	156.4	7.7 ±6.1
Berrien	8	1,728	4.6 ±3.2	11.2	1,782.0	6.3 ±1.6
Branch	5	530	*	4.2	539.6	7.8 ±3.3
Calhoun	15	1,536	9.8 ±4.9	11.0	1,641.6	6.7 ±1.8
Cass	3	507	*	3.4	495.4	6.9 ±3.3
Charlevoix	2	217	*	1.2	226.6	5.3 ±4.2
Cheboygan	2	229	*	0.8	211.8	*
Chippewa	1	324	*	1.2	344.8	3.5 ±2.8
Clare	3	331	*	1.6	326.4	4.9 ±3.4
Clinton	2	824	*	4.0	814.8	4.9 ±2.1
Crawford	1	138	*	1.4	122.4	11.4 ±8.4
Delta	1	345	*	1.6	361.0	4.4 ±3.1
Dickinson	-	247	-	1.4	256.6	5.5 ±4.0
Eaton	3	1,131	*	7.4	1,170.2	6.3 ±2.0
Emmet	1	287	*	1.0	296.8	*
Genesee	37	4,713	7.9 ±2.5	42.0	4,851.2	8.7 ±1.2
Gladwin	-	245	-	2.0	263.0	7.6 ±4.7
Gogebic	-	128	-	1.2	123.8	9.7 ±7.7
Gd. Traverse	3	921	*	4.8	939.8	5.1 ±2.0
Gratiot	2	385	*	2.2	393.8	5.6 ±3.3
Hillsdale	6	534	11.2 ±8.9	4.2	550.6	7.6 ±3.3

Table 3
Number of Infant Deaths, Live Births and Infant Death Rates by
Michigan and Michigan County of Residence,
2017 and 2013 - 2017 Average

County of Residence	2017			2013 - 2017		
	Infant Deaths	Live Births	Infant Death Rate	Average Infant Deaths	Average Live Births	Average Infant Death Rate
Houghton	5	372	*	2.2	367.8	6.0 ±3.5
Huron	3	314	*	0.8	304.2	*
Ingham	16	3,175	5.0 ±2.5	21.4	3,264.4	6.6 ±1.2
Ionia	2	713	*	3.4	729.8	4.7 ±2.2
Iosco	4	228	*	1.8	231.2	7.8 ±5.1
Iron	1	95	*	0.4	92.4	*
Isabella	8	681	11.7 ±8.1	3.6	672.6	5.4 ±2.5
Jackson	14	1,770	7.9 ±4.1	12.8	1,774.4	7.2 ±1.8
Kalamazoo	35	3,055	11.5 ±3.8	21.0	3,151.6	6.7 ±1.3
Kalkaska	-	180	-	1.4	172.4	8.1 ±6.0
Kent	46	8,684	5.3 ±1.5	49.8	8,805.4	5.7 ±0.7
Keweenaw	-	13	-	0.2	17.4	*
Lake	2	123	*	0.6	104.2	*
Lapeer	3	788	*	4.0	807.6	5.0 ±2.2
Leelanau	-	175	-	1.2	172.2	7.0 ±5.6
Lenawee	5	980	*	7.4	1,042.2	7.1 ±2.3
Livingston	8	1,808	4.4 ±3.1	7.8	1,792.4	4.4 ±1.4
Luce	-	60	-	0.8	50.4	*
Mackinac	-	81	-	1.0	87.4	*
Macomb	68	9,252	7.3 ±1.7	59.4	9,365.4	6.3 ±0.7
Manistee	3	210	*	1.2	200.8	6.0 ±4.8
Marquette	1	587	*	2.2	622.0	3.5 ±2.1
Mason	-	301	-	1.0	295.8	*
Mecosta	1	414	*	2.6	418.6	6.2 ±3.4
Menominee	-	179	-	0.6	205.4	*
Midland	2	871	*	3.6	856.8	4.2 ±1.9
Missaukee	-	191	-	0.6	181.4	*
Monroe	11	1,504	7.3 ±4.3	9.0	1,504.6	6.0 ±1.7
Montcalm	3	721	*	3.4	718.4	4.7 ±2.2
Montmorency	-	68	-	-	66.0	-

Table 3
Number of Infant Deaths, Live Births and Infant Death Rates by
Michigan and Michigan County of Residence,
2017 and 2013 - 2017 Average

County of Residence	2017			2013 - 2017		
	Infant Deaths	Live Births	Infant Death Rate	Average Infant Deaths	Average Live Births	Average Infant Death Rate
Muskegon	15	2,125	7.1 ±3.6	17.2	2,111.0	8.1 ±1.7
Newaygo	4	539	*	4.0	545.8	7.3 ±3.2
Oakland	71	13,184	5.4 ±1.2	77.4	13,462.6	5.7 ±0.6
Oceana	-	263	-	2.2	289.8	7.6 ±4.5
Ogemaw	1	204	*	0.6	190.6	*
Ontonagon	-	31	-	0.2	29.8	*
Osceola	2	268	*	1.4	255.6	5.5 ±4.0
Oscoda	-	78	-	0.2	87.4	*
Otsego	1	255	*	1.8	249.6	7.2 ±4.7
Ottawa	11	3,253	3.4 ±2.0	17.4	3,338.4	5.2 ±1.1
Presque Isle	2	101	*	1.2	96.6	12.4 ±9.9
Roscommon	-	163	-	0.6	164.0	*
Saginaw	10	2,248	4.4 ±2.8	19.2	2,246.2	8.5 ±1.7
St. Clair	7	1,462	4.8 ±3.5	8.4	1,543.6	5.4 ±1.6
St. Joseph	4	747	*	4.6	796.6	5.8 ±2.4
Sanilac	3	434	*	3.2	439.0	7.3 ±3.6
Schoolcraft	-	71	-	-	65.0	-
Shiawasee	5	672	*	4.2	708.6	5.9 ±2.5
Tuscola	2	555	*	2.2	556.8	4.0 ±2.3
Van Buren	3	867	*	5.0	875.4	5.7 ±2.2
Washtenaw	15	3,553	4.2 ±2.1	19.6	3,664.6	5.3 ±1.1
Wayne	245	23,257	10.5 ±1.3	221.8	23,365.8	9.5 ±0.6
Wexford	4	374	*	2.6	403.2	6.4 ±3.5
Unknown	-	3	-	-	2.4	-

Note: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2013 - 2017 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 4
Number of Infant Deaths, Live Births and Infant Death Rates for
Selected Cities of Residence, 2017 and 2013 - 2017 Average

City of Residence	2017			2013 - 2017		
	Infant Deaths	Live Births	Infant Death Rate	Average Infant Deaths	Average Live Births	Average Infant Death Rate
Ann Arbor	5	998	*	4.4	1,071.8	4.1 ±1.7
Battle Creek	9	672	13.4 ±8.7	7.0	810.6	8.6 ±2.8
Dearborn	9	1,719	5.2 ±3.4	9.4	1,690.0	5.6 ±1.6
Dearborn Heights	2	827	*	4.4	829.4	5.3 ±2.2
Detroit	137	9,643	14.2 ±2.4	133.6	9,766.0	13.7 ±1.0
East Lansing	-	-	-	-	-	-
Farmington Hills	10	913	11.0 ±6.8	5.8	889.8	6.5 ±2.4
Flint	17	1,397	12.2 ±5.7	18.6	1,562.6	11.9 ±2.4
Grand Rapids	20	3,001	6.7 ±2.9	20.4	3,176.8	6.4 ±1.2
Kalamazoo	9	957	9.4 ±6.1	8.0	1,131.8	7.1 ±2.2
Kentwood	6	796	7.5 ±6.0	5.6	792.4	7.1 ±2.6
Lansing	13	1,728	7.5 ±4.1	16.2	1,818.0	8.9 ±1.9
Livonia	5	869	*	5.0	893.4	5.6 ±2.2
Midland	1	441	*	2.6	463.8	5.6 ±3.0
Muskegon	3	516	*	7.8	585.2	13.3 ±4.2
Novi	2	619	*	3.2	643.8	5.0 ±2.4
Pontiac	11	1,072	10.3 ±6.0	13.6	1,106.2	12.3 ±2.9
Portage	9	550	16.4 ±10.6	3.8	567.8	6.7 ±3.0
Rochester Hills	3	727	*	2.6	749.0	3.5 ±1.9
Roseville	3	569	*	4.8	598.4	8.0 ±3.2
Royal Oak	3	716	*	2.6	746.0	3.5 ±1.9
Saginaw	6	788	7.6 ±6.1	11.6	821.2	14.1 ±3.6
Southfield	5	769	*	6.8	784.6	8.7 ±2.9
St Clair Shores	2	575	*	3.2	583.2	5.5 ±2.7
Sterling Heights	9	1,436	6.3 ±4.1	9.6	1,440.0	6.7 ±1.9
Taylor	9	809	11.1 ±7.2	6.0	855.0	7.0 ±2.5
Troy	4	822	*	3.6	813.8	4.4 ±2.0
Warren	14	1,700	8.2 ±4.3	10.8	1,689.0	6.4 ±1.7
Westland	8	1,058	7.6 ±5.2	7.8	1,080.6	7.2 ±2.3
Wyoming	5	1,171	*	7.0	1,202.0	5.8 ±1.9

Note: This table shows cities with (1) populations greater than 40,000 according to 2007 estimates and (2) three-year average births of greater than 200. City birth and mortality statistics have changed from pre-2011 reports due to geocoding. See Data Quality Considerations for further details. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2013-2017 Geocoded Michigan Death Certificate Registries 2013-2017 Geocoded Michigan Birth Certificate Registries Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 5
Number and Rate of Infant Deaths by Race for
Residents of Michigan, Detroit City, Wayne County (Excluding Detroit City)
and Michigan (Excluding Wayne County), 1990- 2017

Year	Race	Michigan		Detroit City		Wayne County (excluding Detroit)		Michigan (excluding Wayne County)	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
1990	Total	1,638	10.7 ±0.5	491	20.3 ±1.8	170	10.3 ±1.5	977	8.7 ±0.5
	White	930	7.9 ±0.5	36	8.7 ±2.8	130	8.8 ±1.5	764	7.7 ±0.5
	Black	689	21.6 ±1.6	451	23.0 ±2.1	38	24.8 ±7.8	200	18.7 ±2.6
1991	Total	1,554	10.4 ±0.5	456	19.4 ±1.8	139	8.4 ±1.4	959	8.8 ±0.6
	White	857	7.5 ±0.5	39	10.0 ±3.1	103	7.1 ±1.4	715	7.4 ±0.5
	Black	683	21.7 ±1.6	417	21.6 ±2.1	34	20.3 ±6.7	232	21.9 ±2.8
1992	Total	1,460	10.2 ±0.5	462	21.4 ±1.9	128	8.1 ±1.4	870	8.2 ±0.5
	White	781	7.0 ±0.5	43	12.3 ±3.7	92	6.6 ±1.4	646	6.9 ±0.5
	Black	656	22.1 ±1.7	415	23.3 ±2.2	34	21.8 ±7.2	207	20.1 ±2.7
1993	Total	1,319	9.5 ±0.5	348	17.2 ±1.8	111	7.1 ±1.3	860	8.3 ±0.6
	White	770	7.1 ±0.5	28	9.1 ±3.3	88	6.4 ±1.3	654	7.2 ±0.5
	Black	530	18.8 ±1.6	317	18.7 ±2.0	21	14.4 ±6.1	192	19.4 ±2.7
1994	Total	1,184	8.6 ±0.5	322	16.6 ±1.8	105	6.8 ±1.3	757	7.3 ±0.5
	White	670	6.3 ±0.5	28	9.4 ±3.5	69	5.1 ±1.2	573	6.3 ±0.5
	Black	495	18.2 ±1.6	293	18.2 ±2.1	35	23.7 ±7.7	167	17.5 ±2.6
1995	Total	1,110	8.3 ±0.5	274	15.6 ±1.8	89	5.8 ±1.2	747	7.4 ±0.5
	White	655	6.2 ±0.5	17	6.0 ±2.8	64	4.8 ±1.2	574	6.4 ±0.5
	Black	432	17.3 ±1.6	254	17.6 ±2.1	25	18.2 ±7.1	153	16.8 ±2.6
1996	Total	1,072	8.0 ±0.5	248	15.0 ±1.8	101	6.7 ±1.3	723	7.1 ±0.5
	White	627	6.0 ±0.5	9	3.4 ±2.2	74	5.6 ±1.3	544	6.1 ±0.5
	Black	421	17.5 ±1.7	238	17.5 ±2.2	23	16.2 ±6.6	160	17.7 ±2.7
1997	Total	1,085	8.1 ±0.5	253	15.1 ±1.8	100	6.6 ±1.3	732	7.2 ±0.5
	White	640	6.1 ±0.5	23	8.6 ±3.5	76	5.8 ±1.3	541	6.1 ±0.5
	Black	425	17.6 ±1.7	229	16.7 ±2.1	23	16.5 ±6.7	173	19.1 ±2.8

Table 5
Number and Rate of Infant Deaths by Race for
Residents of Michigan, Detroit City, Wayne County (Excluding Detroit City)
and Michigan (Excluding Wayne County), 1990- 2017

Year	Race	Michigan		Detroit City		Wayne County (excluding Detroit)		Michigan (excluding Wayne County)	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
1998	Total	1,091	8.2 ±0.5	241	14.7 ±1.8	99	6.6 ±1.3	751	7.4 ±0.5
	White	660	6.3 ±0.5	15	5.8 ±2.9	74	5.7 ±1.3	571	6.4 ±0.5
	Black	405	16.8 ±1.6	219	16.1 ±2.1	22	14.9 ±6.2	164	18.0 ±2.7
1999	Total	1,071	8.0 ±0.5	245	15.5 ±1.9	97	6.4 ±1.3	729	7.1 ±0.5
	White	616	5.9 ±0.5	15	5.6 ±2.8	72	5.6 ±1.3	529	5.9 ±0.5
	Black	428	17.9 ±1.7	228	17.8 ±2.3	23	15.1 ±6.1	177	18.6 ±2.7
2000	Total	1,112	8.2 ±0.5	232	14.7 ±1.9	98	6.4 ±1.3	782	7.4 ±0.5
	White	637	6.0 ±0.5	20	7.5 ±3.3	64	5.0 ±1.2	553	6.1 ±0.5
	Black	437	18.2 ±1.7	206	16.1 ±2.2	33	20.2 ±6.8	198	20.5 ±2.8
2001	Total	1,066	8.0 ±0.5	210	13.9 ±1.9	81	5.3 ±1.2	775	7.5 ±0.5
	White	635	6.1 ±0.5	14	5.3 ±2.8	58	4.5 ±1.2	563	6.3 ±0.5
	Black	398	16.9 ±1.7	195	15.9 ±2.2	21	12.9 ±5.5	182	18.9 ±2.7
2002	Total	1,054	8.1 ±0.5	239	16.8 ±2.1	103	6.9 ±1.3	712	7.1 ±0.5
	White	611	6.0 ±0.5	20	8.0 ±3.5	70	5.7 ±1.3	521	6.0 ±0.5
	Black	410	18.4 ±1.8	217	19.1 ±2.5	28	16.9 ±6.2	165	17.9 ±2.7
2003	Total	1,112	8.5 ±0.5	227	16.2 ±2.1	120	8.1 ±1.4	765	7.5 ±0.5
	White	680	6.7 ±0.5	28	11.7 ±4.3	85	7.1 ±1.5	567	6.5 ±0.5
	Black	391	17.5 ±1.7	199	17.9 ±2.5	29	16.9 ±6.1	163	17.1 ±2.6
2004	Total	984	7.6 ±0.5	211	15.6 ±2.1	89	6.1 ±1.3	684	6.7 ±0.5
	White	527	5.2 ±0.4	12	5.0 ±2.8	54	4.6 ±1.2	461	5.3 ±0.5
	Black	388	17.3 ±1.7	193	17.9 ±2.5	30	16.1 ±5.7	165	16.8 ±2.5
2005	Total	1,013	7.9 ±0.5	210	16.0 ±2.1	93	6.5 ±1.3	710	7.1 ±0.5
	White	549	5.5 ±0.5	13	5.7 ±3.1	58	5.1 ±1.3	478	5.6 ±0.5
	Black	400	17.9 ±1.7	188	17.9 ±2.5	34	17.7 ±5.9	178	17.9 ±2.6

Table 5
Number and Rate of Infant Deaths by Race for
Residents of Michigan, Detroit City, Wayne County (Excluding Detroit City)
and Michigan (Excluding Wayne County), 1990- 2017

Year	Race	Michigan		Detroit City		Wayne County (excluding Detroit)		Michigan (excluding Wayne County)	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
2006	Total	940	7.4 ±0.5	175	13.5 ±2.0	98	6.8 ±1.4	667	6.7 ±0.5
	White	534	5.4 ±0.5	9	4.0 ±2.6	61	5.5 ±1.4	464	5.4 ±0.5
	Black	339	14.8 ±1.6	153	14.7 ±2.3	31	14.4 ±5.0	155	15.0 ±2.3
2007	Total	997	8.0 ±0.5	179	14.8 ±2.1	95	6.7 ±1.3	723	7.3 ±0.5
	White	549	5.8 ±0.5	13	7.1 ±3.8	59	5.5 ±1.4	477	5.9 ±0.5
	Black	368	16.5 ±1.7	155	16.1 ±2.5	25	11.4 ±4.5	188	17.9 ±2.5
2008	Total	894	7.4 ±0.5	175	14.8 ±2.2	100	7.2 ±1.4	619	6.5 ±0.5
	White	496	5.4 ±0.5	8	4.7 ±3.3	63	5.9 ±1.5	425	5.4 ±0.5
	Black	332	14.6 ±1.6	159	16.7 ±2.6	27	12.3 ±4.6	146	13.3 ±2.1
2009	Total	881	7.5 ±0.5	166	14.8 ±2.2	82	6.1 ±1.3	633	6.8 ±0.5
	White	470	5.4 ±0.5	6	4.3 ±3.5	54	5.4 ±1.4	410	5.4 ±0.5
	Black	346	15.5 ±1.6	152	16.7 ±2.6	19	7.9 ±3.5	175	16.2 ±2.4
2010	Total	817	7.1 ±0.5	146	13.3 ±2.2	87	6.6 ±1.4	584	6.4 ±0.5
	White	462	5.5 ±0.5	10	9.6 ±5.9	44	4.6 ±1.4	408	5.5 ±0.5
	Black	312	14.2 ±1.6	128	14.3 ±2.5	37	15.0 ±4.8	147	13.8 ±2.2
2011	Total	749	6.6 ±0.5	130	12.6 ±2.1	100	7.5 ±1.5	519	5.7 ±0.5
	White	411	4.9 ±0.5	5	*	58	6.0 ±1.5	348	4.7 ±0.5
	Black	300	13.7 ±1.5	119	14.0 ±2.5	37	14.6 ±4.7	144	13.2 ±2.1
2012	Total	783	6.9 ±0.5	151	15.0 ±2.4	87	6.7 ±1.4	545	6.1 ±0.5
	White	455	5.5 ±0.5	12	15.7 ±8.8	56	5.9 ±1.6	387	5.3 ±0.5
	Black	291	13.5 ±1.5	133	16.0 ±2.7	27	11.2 ±4.2	131	12.0 ±2.0
2013	Total	799	7.0 ±0.5	132	13.1 ±2.2	84	6.2 ±1.3	583	6.5 ±0.5
	White	473	5.7 ±0.5	7	8.8 ±6.5	57	6.0 ±1.6	409	5.6 ±0.5
	Black	290	13.1 ±1.5	119	14.4 ±2.6	22	8.2 ±3.4	149	13.5 ±2.1

Table 5
Number and Rate of Infant Deaths by Race for
Residents of Michigan, Detroit City, Wayne County (Excluding Detroit City)
and Michigan (Excluding Wayne County), 1990- 2017

Year	Race	Michigan		Detroit City		Wayne County (excluding Detroit)		Michigan (excluding Wayne County)	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
2014	Total	773	6.8 ±0.5	130	13.2 ±2.3	88	6.5 ±1.4	555	6.1 ±0.5
	White	452	5.3 ±0.5	8	9.6 ±6.6	55	5.9 ±1.6	389	5.2 ±0.5
	Black	284	13.3 ±1.5	120	14.7 ±2.6	25	9.7 ±3.8	139	13.2 ±2.2
2015	Total	770	6.8 ±0.5	142	14.4 ±2.3	83	6.1 ±1.3	545	6.1 ±0.5
	White	423	5.1 ±0.5	9	10.1 ±6.6	60	6.4 ±1.6	354	4.9 ±0.5
	Black	311	14.5 ±1.6	125	15.4 ±2.7	17	6.4 ±3.1	169	15.8 ±2.4
2016	Total	747	6.6 ±0.5	125	13.3 ±2.3	80	5.8 ±1.3	542	6.0 ±0.5
	White	401	4.8 ±0.5	7	8.0 ±5.9	49	5.1 ±1.4	345	4.7 ±0.5
	Black	294	13.7 ±1.6	105	13.6 ±2.6	30	10.8 ±3.8	159	14.4 ±2.2
2017	Total	762	6.8 ±0.5	137	14.2 ±2.4	108	7.9 ±1.5	517	5.9 ±0.5
	White	392	4.8 ±0.5	7	7.2 ±5.3	45	4.7 ±1.4	340	4.8 ±0.5
	Black	320	14.6 ±1.6	122	15.5 ±2.7	52	18.8 ±5.1	146	13.0 ±2.1

Note: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Source: 1990- 2017 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 6
Number and Rate of Infant Deaths by Race,
Michigan and United States Residents, 1989 - 2017

Year	United States						Michigan					
	All Races		White		Black		All Races		White		Black	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
1989	39,655	9.8	25,794	8.0	12,527	18.6	1,645	11.1	946	8.2	672	22.5
1990	38,351	9.2	24,883	7.6	12,290	18.0	1,638	10.7	930	7.9	689	21.6
1991	36,766	8.9	23,657	7.3	11,994	17.6	1,554	10.4	857	7.5	683	21.7
1992	34,628	8.5	22,164	6.9	11,348	16.8	1,460	10.2	781	7.0	656	22.1
1993	33,466	8.4	21,497	6.8	10,887	16.5	1,319	9.5	770	7.1	530	18.8
1994	31,710	8.0	20,504	6.6	10,072	15.8	1,184	8.6	670	6.3	495	18.2
1995	29,583	7.6	19,490	6.3	9,118	15.1	1,110	8.3	655	6.2	432	17.3
1996	28,487	7.3	18,761	6.1	8,730	14.7	1,072	8.0	627	6.0	421	17.5
1997	28,045	7.2	18,539	6.0	8,496	14.2	1,085	8.1	640	6.1	425	17.6
1998	28,371	7.2	18,561	6.0	8,726	14.3	1,091	8.2	660	6.3	405	16.8
1999	27,953	7.1	18,069	5.8	8,832	14.6	1,071	8.0	616	5.9	428	17.9
2000	28,035	6.9	18,144	5.7	8,771	14.0	1,112	8.2	637	6.0	437	18.2
2001	27,568	6.8	17,955	5.7	8,498	14.0	1,066	8.0	635	6.1	398	16.9
2002	28,034	7.0	18,369	5.8	8,524	14.4	1,054	8.1	611	6.0	410	18.4
2003	28,025	6.9	18,440	5.7	8,402	14.0	1,112	8.5	680	6.7	391	17.5
2004	27,936	6.8	18,231	5.7	8,494	13.3	984	7.6	527	5.2	388	17.3
2005	28,440	6.9	18,514	5.7	8,695	13.7	1,013	7.9	549	5.5	400	17.9
2006	28,527	6.7	18,403	5.6	8,858	13.3	940	7.4	534	5.4	339	14.8
2007	29,138	6.8	18,807	5.6	8,944	13.2	997	8.0	549	5.8	368	16.5
2008	28,059	6.6	18,164	5.6	8,543	12.7	894	7.4	496	5.4	332	14.6
2009	26,412	6.4	16,817	5.3	8,312	12.6	881	7.5	470	5.4	346	15.5
2010	24,548	6.1	15,933	5.2	7,388	11.6	817	7.1	462	5.5	312	14.2
2011	23,910	6.1	15,453	5.1	7,058	11.2	749	6.6	411	4.9	300	13.7
2012	23,654	6.0	15,264	5.1	6,896	10.9	783	6.9	455	5.5	291	13.5
2013	23,440	6.0	15,125	5.0	7,123	11.2	799	7.0	473	5.7	290	13.1
2014	23,215	5.8	14,883	5.1	7,076	10.4	773	6.8	452	5.3	284	13.3
2015	23,455	5.9	14,834	5.0	7,289	10.6	770	6.8	423	5.1	311	14.5
2016	23,161	5.9	14,568	4.9	7,229	10.6	747	6.6	401	4.8	294	13.7
2017	---	---	---	---	---	---	762	6.8	392	4.8	320	14.6

Note: Rates are per 1,000 live births.

Source: 1989 - 2017 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.
 2007-2012 NCHS, Linked Birth / Infant Death Records 2007-2015 on CDC WONDER On-line Database. Accessed at <http://wonder.cdc.gov/lbd-current.html> on Mar 7, 2017. "Deaths: Final Data for 2006", Vol 57, No. 14, April 2009; and [National Center for Health Statistics, historical data.](#)

Table 7-A
Number of Hispanic and Middle Easterner Infant Deaths by Selected County and Michigan
Michigan Residents, 2007 - 2017

Hispanic Infant Deaths

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Michigan	90	74	74	57	44	74	80	61	77	68	58
Hispanic Infant Death Rates	10.3	8.5	9.0	7.0	5.6	9.7	10.3	8.0	9.9	8.8	7.6
Allegan	-	2	1	-	1	-	3	-	-	1	2
Genesee	5	1	1	1	2	-	-	1	2	4	3
Ingham	3	2	8	1	1	2	4	2	1	2	2
Kalamazoo	4	3	-	4	-	2	2	1	4	1	1
Kent	13	16	6	8	12	10	6	7	13	10	4
Lenawee	2	-	2	1	1	2	5	2	-	1	1
Macomb	6	2	4	-	-	1	3	-	5	2	3
Muskegon	2	2	3	-	1	1	5	-	2	2	1
Oakland	5	9	9	6	7	8	8	4	9	1	5
Ottawa	3	3	4	1	2	4	8	2	4	4	3
Saginaw	8	4	5	2	2	5	4	4	2	8	-
Van Buren	2	2	-	1	-	2	1	2	3	1	-
Washtenaw	-	3	4	2	-	2	1	-	-	1	-
Wayne	19	15	13	15	8	16	17	13	21	13	18
All Other Counties	18	10	14	15	7	19	13	23	11	17	15

Table 7-B
Number of Hispanic and Middle Easterner Infant Deaths by Selected County and Michigan
Michigan Residents, 2007 - 2017 (Continued)

Middle Easterner Infant Deaths

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Michigan	32	34	22	24	24	26	36	21	26	30	22
Arabic Infant Death Rates	8.0	8.5	5.8	5.7	5.9	6.2	7.8	4.4	5.3	6.0	4.1
Macomb	3	7	6	5	2	5	6	3	2	6	4
Oakland	4	8	2	6	3	5	7	3	4	5	2
Wayne	23	16	10	10	15	14	17	9	16	13	14
All Other Counties	2	3	4	3	4	2	6	6	4	6	2

Note: The increase in hispanic mortality incidence reflects a change in death certificate reporting. Michigan's death certificate was revised in 2003 to comply with a change in the National Standard death certificate and a separate indicator of hispanic ancestry was introduced. In 2005, the hispanic ancestry indicator was used in tabulations for the first time, resulting in the identification of hispanic deaths that might otherwise have been missed. The increase in hispanic death cases between 2004 and 2005 should not be considered as a change in trend.

Caution: Care should be taken drawing inferences from rates based on small numbers of events or small population base. These rates tend to exhibit considerable variation which may negate their usefulness for comparative purpose

Source: 2017 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.


Table 8
Three-year Average Infant Death Rates by Race and Ancestry,
Michigan Residents, 1999 - 2017

Trends	Race				Ancestry	
	White	Black	Native American	Asian and Pacific Islander	Hispanic	Middle Easterner
1999-2001	6.0	17.7	10.5	4.3	7.3	5.9
2000-2002	6.0	17.8	11.6	4.2	7.4	5.4
2001-2003	6.2	17.6	12.4	4.2	8.3	5.2
2002-2004	6.0	17.7	10.0	4.7	9.2	4.7
2003-2005	5.8	17.5	10.1	4.5	10.1	4.9
2004-2006	5.4	16.6	8.8	3.8	10.6	5.8
2005-2007	5.6	16.4	10.6	4.0	10.8	7.3
2006-2008	5.6	15.3	9.9	4.2	9.9	8.1
2007-2009	5.6	15.5	9.3	4.8	9.3	7.5
2008-2010	5.4	14.8	9.1	3.8	8.2	6.7
2009-2011	5.2	14.5	9.6	3.8	7.2	5.8
2010-2012	5.3	13.8	11.2	3.6	7.4	5.9
2011-2013	5.3	13.4	10.1	4.1	8.5	6.7
2012-2014	5.5	13.3	11.9	4.4	9.4	6.1
2013-2015	5.4	13.6	10.2	4.1	9.1	5.8
2014-2016	5.1	13.8	14.0	3.7	8.6	5.2
2015-2017	4.9	14.3	14.2	3.5	8.4	5.1

Note: Rates are per 1,000 live births.

Source: 1970 - 2017 Michigan Resident Birth and Death Files, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Figure 3
Three-year Average Infant Mortality Rates by Race and Ancestry,
Michigan Residents, 1998-2017



Note: Rates are infant deaths per 1,000 live births.

Source: 1980-2017 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 9
Infant, Hebdomadal, Fetal and Perinatal Death Rates
by Specified Race and Ancestry
Michigan Residents, 2017

Race/Ancestry	Live Births	Infant Deaths		Hebdomadal Deaths		Fetal Deaths		Perinatal Deaths	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
All Races	111,507	762	6.8 ±0.5	441	4.0 ±0.4	605	5.4 ±0.4	1,046	9.3 ±0.6
White	80,967	392	4.8 ±0.5	232	2.9 ±0.4	328	4.0 ±0.4	560	6.9 ±0.6
Black	21,897	320	14.6 ±1.6	174	7.9 ±1.2	222	10.0 ±1.3	396	17.9 ±1.7
American Indian	643	10	15.6 ±9.6	3	*	5	*	8	12.3 ±8.5
Asian and Pacific Islander	4,675	19	4.1 ±1.8	15	3.2 ±1.6	28	6.0 ±2.2	43	9.1 ±2.7
Middle Easterner	5,315	22	4.1 ±1.7	11	2.1 ±1.2	15	2.8 ±1.4	26	4.9 ±1.9
Hispanic	7,639	58	7.6 ±1.9	39	5.1 ±1.6	38	4.9 ±1.6	77	10.0 ±2.2

Note: Hebdomadal deaths are deaths to infants ages zero to six days. Perinatal deaths are hebdomadal plus fetal deaths.

Infant death and hebdomadal rates are per 1,000 live births. Fetal death and perinatal death rates are per 1,000 live births plus fetal deaths.

Rates are calculated using live births by race/ancestry of mother as denominator. Live births are by the race/ancestry of the mother on the birth certificate. Infant and hebdomadal deaths are by the race on the death certificate. Fetal deaths are by the race of the mother on the fetal death certificate. Perinatal deaths are by mother's race/ancestry on the fetal death certificate or the infant's race/ancestry on the death certificate.

Asterisk (*) indicates that data do not meet standards of precision or reliability.

Caution: Care should be taken drawing inferences from rates based on small numbers of events or small population base. These rates tend to exhibit considerable variation which may negate their usefulness for comparative purposes.

Source: 2017 Michigan Resident Infant Death File, 2017 Michigan Fetal Death Registry
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 10
Infant, Hebdomadal, Fetal and Perinatal Death Rates
Michigan Residents, 2007-2017

Year	Live Births	Infant Deaths		Hebdomadal Deaths		Fetal Deaths		Perinatal Deaths	
		Number	Rate	Number	Rate	Number	Rate	Number	Rate
2007	125,172	997	8.0	553	4.4	731	5.8	1,284	10.2
2008	121,231	894	7.4	501	4.1	762	6.2	1,263	10.4
2009	117,309	881	7.5	478	4.1	692	5.9	1,170	9.9
2010	114,717	817	7.1	475	4.1	650	5.6	1,125	9.8
2011	114,159	749	6.6	414	3.6	633	5.5	1,047	9.1
2012	112,708	783	6.9	459	4.1	591	5.2	1,050	9.3
2013	113,732	799	7.0	431	3.8	580	5.1	1,011	8.8
2014	114,460	773	6.8	417	3.6	625	5.4	1,042	9.1
2015	113,211	770	6.8	400	3.5	547	4.8	941	8.4
2016	113,374	747	6.6	393	3.5	618	5.4	1,011	8.9
2017	111,507	762	6.8	441	4.0	605	5.4	1,046	9.3

Note: Hebdomadal deaths are deaths to infants ages zero to six days. Perinatal deaths are hebdomadal plus fetal deaths.

Infant death and hebdomadal rates are per 1,000 live births. Fetal death and perinatal death rates are per 1,000 live births plus fetal deaths.

In 2017, hospital audits found additional fetal deaths. The number of fetal deaths between 2007-2015 will not match what was reported in the past.

Source: 2017 Michigan Resident Infant Death File, 2017 Michigan Fetal Death Registry
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 11
Number of Infant Deaths, Live Births and Infant Death Rates
by Selected Characteristics of Newborn and of Mother
for Michigan Residents, 2017

Characteristics of Newborn	Infant Deaths	Live Births	Infant Death Rate
Sex of Infant	762	111,507	6.8 ±0.5
Male	427	56,906	7.5 ±0.7
Female	332	54,560	6.1 ±0.7
Birthweight	762	111,507	6.8 ±0.5
1,499 Grams or Less	416	1,687	246.6 ±20.9
1,500 - 2,499 Grams	109	8,158	13.4 ±2.5
2,500 Grams or More	235	101,599	2.3 ±0.3
Plurality	762	111,507	6.8 ±0.5
Single Birth	639	107,166	6.0 ±0.5
Multiple Birth	122	4,302	28.4 ±5.0
Characteristics of Mother	Infant Deaths	Live Births	Infant Death Rate
Age of Mother	762	111,507	6.8 ±0.5
< 20	48	5,363	9.0 ±2.5
20 - 29	424	58,631	7.2 ±0.7
30 - 39	268	44,575	6.0 ±0.7
40 +	20	2,897	6.9 ±3.0
Marital Status	762	111,507	6.8 ±0.5
Married	310	65,288	4.7 ±0.5
Unmarried	451	46,180	9.8 ±0.9
Level of Prenatal Care	762	111,507	6.8 ±0.5
Adequate	390	75,429	5.2 ±0.5
Intermediate	201	25,515	7.9 ±1.1
Inadequate	169	10,509	16.1 ±2.4

Table 11
Number of Infant Deaths, Live Births and Infant Death Rates
by Selected Characteristics of Newborn and of Mother
for Michigan Residents, 2017

Exposure to Tobacco	Infant Deaths	Live Births	Infant Death Rate
Smoked During Pregnancy	762	111,507	6.8 ±0.5
No	606	95,416	6.4 ±0.5
Yes	156	16,052	9.7 ±1.5
Secondhand Smoke During Pregnancy	762	111,507	6.8 ±0.5
No	581	94,275	6.2 ±0.5
Yes	114	13,448	8.5 ±1.5
Smoked or Secondhand Smoke During Pregnancy	762	111,507	6.8 ±0.5
No	508	86,827	5.9 ±0.5
Yes	196	21,535	9.1 ±1.3
History of smoking	762	111,507	6.8 ±0.5
No	560	91,262	6.1 ±0.5
Yes	174	19,716	8.8 ±1.3

Note: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence. Records with selected characteristics unknown are included in total only. Level of prenatal care is according to the Kessner Index which is a classification of prenatal care based on the month of pregnancy in which prenatal care began, the number of prenatal visits and the length of pregnancy (i.e. for shorter pregnancies, fewer prenatal visits constitute adequate care.)

Source: 2017 Michigan Resident Birth-Death Matched File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 12
Number of Infant Deaths by Cause of Death,
1980-2017, Michigan Residents

Causes of Death	1980	1985	1990	1995	2000	2005	2010	2012	2013	2014	2015	2016	2017
All Causes	2,280	1,575	1,638	1,110	1,112	1,013	817	783	799	773	770	747	762
Certain gastrointestinal diseases	7	6	6	6	10	12	9	7	5	6	7	3	5
Septicemia	8	6	17	2	5	10	4	8	2	8	3	4	4
Remainder of infectious and parasitic diseases	5	11	18	12	7	6	3	2	3	5	4	4	3
Meningitis	13	7	11	2	3	3	5	2	1	1	-	3	1
Pneumonia and Influenza	27	11	26	20	20	6	8	5	7	7	5	2	9
Congenital anomalies	321	330	300	204	192	197	184	127	160	139	152	149	141
Newborn affected by maternal complications of pregnancy	72	52	108	47	45	67	40	68	52	42	45	36	31
Newborn affected by complications of placenta, cord and membranes	50	34	24	33	36	28	26	28	30	28	31	28	30
Disorders relating to short gestation and unspecified low birthweight	145	150	251	219	262	219	198	188	176	174	149	161	196
Birth trauma	45	14	16	8	2	1	-	-	-	-	-	1	-
Intrauterine hypoxia and birth asphyxia	69	43	25	11	24	19	20	14	8	7	11	5	15
Respiratory distress syndrome	206	176	164	82	65	57	23	26	20	19	14	18	22
Other respiratory conditions	199	132	120	63	65	52	27	27	20	22	17	21	13
Infections specific to the perinatal period	35	32	33	23	35	42	27	21	22	24	21	23	28
Neonatal hemorrhage	38	10	8	13	28	23	12	12	9	18	16	9	9
All other and ill-define conditions originating in the perinatal period	169	168	73	47	39	46	27	34	47	28	39	31	24
Sudden infant death syndrome	260	228	257	148	99	47	44	15	31	35	33	45	32
Accidents and adverse effects	39	35	38	40	55	68	66	93	90	89	97	82	82
Homicides	7	8	17	12	13	15	13	14	5	13	6	9	14
All other causes	136	122	126	118	107	95	81	92	111	108	120	113	103

Note: Causes of death are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. See [Selected Causes of Groups for Infant Mortality](#) for details of grouping and comparability.

Source: 1980-2017 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 13
Three-year Moving Average Infant Deaths Rates per 10,000 Live Births, by Cause of Death,
1980-2017, Michigan Residents


Causes of Death	1980 to 1982	1985 to 1987	1990 to 1992	1995 to 1997	2000 to 2002	2005 to 2007	2010 to 2012	2011 to 2013	2012 to 2014	2013 to 2015	2014 to 2016	2015 to 2017
Total Infant Deaths	126.8	112.4	104.2	81.5	81.0	77.6	68.8	68.4	69.1	68.6	67.1	67.4
Certain gastrointestinal diseases	.6	.6	.6	.6	.6	.9	.8	.7	.5	.5	.5	.4
Septicemia	.4	.5	.9	.4	.5	.6	.4	.3	.5	.4	.4	.3
Remainder of infectious and parasitic diseases	.4	.8	1.1	.6	.5	.5	.2	.2	.3	.4	.4	.3
Meningitis	.9	.4	.4	*	*	.3	.2	*	*	*	*	*
Pneumonia and Influenza	1.4	1.1	1.8	1.1	1.1	.5	.6	.5	.6	.6	.4	.5
Congenital anomalies	22.7	21.4	18.1	15.1	13.8	15.1	12.8	12.1	12.5	13.2	12.9	13.1
Newborn affected by maternal complications of pregnancy	5.8	4.6	6.6	2.9	5.7	5.4	4.6	5.0	4.8	4.1	3.6	3.3
Newborn affected by complications of placenta, cord and membranes	3.2	2.6	2.1	2.9	2.4	2.1	2.1	2.2	2.5	2.6	2.6	2.6
Disorders relating to short gestation and unspecified low birthweight	10.6	12.2	18.2	16.0	17.6	17.2	16.8	16.2	15.8	14.6	14.2	15.0
Birth trauma	2.5	.7	.7	.4	.2	*	*	*	-	-	*	*
Intrauterine hypoxia and birth asphyxia	4.5	2.4	1.6	1.2	1.6	.9	1.2	.9	.9	.8	.7	.9
Respiratory distress syndrome	14.8	11.2	9.7	6.1	4.5	3.6	1.8	1.8	1.9	1.6	1.5	1.6
Other respiratory conditions	12.2	10.2	7.1	5.1	4.3	3.5	2.4	2.2	2.0	1.7	1.8	1.5
Infections specific to the perinatal period	2.2	2.3	2.2	2.1	3.0	2.9	2.0	1.9	2.0	2.0	2.0	2.1
Neonatal hemorrhage	2.0	.7	.6	.8	1.3	1.3	1.0	.9	1.1	1.3	1.3	1.0
All other and ill-define conditions originating in the perinatal period	13.3	12.1	5.1	3.7	3.1	4.3	2.9	3.5	3.2	3.3	2.9	2.8
Sudden infant death syndrome	16.6	17.1	16.6	10.8	4.3	3.8	2.7	2.3	2.4	2.9	3.3	3.3
Accidents and adverse effects	2.8	2.1	2.4	2.7	4.6	5.7	7.1	7.8	8.0	8.1	7.9	7.7
Homicides	.4	.9	.9	.9	.8	.8	1.1	.9	.9	.7	.8	.9
All other causes	9.7	8.6	7.6	7.9	8.3	8.3	7.9	8.8	9.1	9.9	10.0	9.9

Note: For purposes of comparing small values, these rates are calculated at deaths per **10,000** live births; in most Michigan Community Health infant death reports, infant deaths are per **1,000**. **Causes of death** are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. See [Selected Causes of Groups for Infant Mortality](#) for details of grouping and comparability.

An asterisk (*) indicates the rate does not meet standards of reliability or precision. A dash (-) indicates quantity zero.

Source: 1980-2017 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Figure 4
Infant Mortality Rates by Leading Causes of Infant Deaths,
Michigan Residents, 1980-2017



Note: For purposes of comparing small values, these rates are calculated at deaths per **10,000** live births; in most Michigan Community Health infant death reports, infant deaths are per **1,000**.

Causes of death are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. See [Selected Causes of Groups for Infant Mortality](#) for details of grouping and comparability.

An asterisk (*) indicates the rate does not meet standards of reliability or precision. A dash (-) indicates quantity zero.

Source: 1980-2017 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

**Table 14-A
Number of Infant Deaths by Cause of Death,
2014-2017, Michigan Residents**

Dollfus Primary Cause of Death Group

Cause of Death Group	Number of Deaths				Percent of Deaths			
	2014	2015	2016	2017	2014	2015	2016	2017
Total	773	770	747	762	100.0	100.0	100.0	100.0
Prematurity & related conditions	247	223	228	255	32.0	29.0	30.5	33.5
Congenital anomalies	139	152	151	141	18.0	19.7	20.2	18.5
Sudden Infant Death Syndrome (SIDS)	35	33	45	32	4.5	4.3	6.0	4.2
Obstetric conditions	47	52	39	38	6.1	6.8	5.2	5.0
Birth asphyxia	7	11	5	15	0.9	1.4	0.7	2.0
Perinatal infections	40	36	37	46	5.2	4.7	5.0	6.0
Other infections	29	25	17	22	3.8	3.2	2.3	2.9
External causes	105	104	98	103	13.6	13.5	13.1	13.5
Other causes	124	134	127	110	16.0	17.4	17.0	14.4

Table 14-A Dollfus Subcategory Cause of Death Group

Dollfus Subcategory	Number of Deaths				Percent of Deaths			
	2014	2015	2016	2017	2014	2015	2016	2017
Total	773	770	747	762	100.0	100.0	100.0	100.0
Short gestation & low birthweight	174	149	161	196	22.5	19.4	21.6	25.7
Birth trauma, neonatal hemorrhage, & hematological disorders	18	15	10	8	2.3	1.9	1.3	1.0
RDS, interstitial emphysema, pulmonary hemorrhage, & atelectasis	35	26	29	29	4.5	3.4	3.9	3.8
Chronic respiratory distress syndrome, & respiratory diseases originating in the perinatal period	1	3	5	4	0.1	0.4	0.7	0.5
Necrotizing enterocolitis	11	17	14	11	1.4	2.2	1.9	1.4
Neonatal cardiac failure & other cardiovascular disorders originating in the perinatal period	8	13	9	7	1.0	1.7	1.2	0.9
Congenital malformations, deformations, & chromosomal anomalies, congenital disorders	139	152	151	141	18.0	19.7	20.2	18.5
Sudden Infant Death Syndrome	35	33	45	32	4.5	4.3	6.0	4.2
Incompetent cervix, premature rupture of membranes	20	26	28	26	2.6	3.4	3.7	3.4
Multiple pregnancy	13	12	3	–	1.7	1.6	0.4	–
Placental previa, separation and placental abnormalities	14	14	8	12	1.8	1.8	1.1	1.6
Intrauterine hypoxia, birth asphyxia	7	11	5	15	0.9	1.4	0.7	2.0
Streptococcal meningitis	1	–	1	1	0.1	–	0.1	0.1
Maternal renal and urinary tract diseases, maternal infections, and parasitic diseases	1	–	–	1	0.1	–	–	0.1
Chorioamnionitis	12	13	10	15	1.6	1.7	1.3	2.0
Bacterial Sepsis, congenital pneumonia, and other infections specific to the perinatal period	26	23	26	29	3.4	3.0	3.5	3.8
Other infectious and parasitic diseases	18	13	11	12	2.3	1.7	1.5	1.6
Bacterial haemophilus and pneumococcal meningitis	1	–	2	–	0.1	–	0.3	–
Influenza and pneumonia and other respiratory infections	10	12	4	10	1.3	1.6	0.5	1.3
Accidents involving transportation	2	2	3	5	0.3	0.3	0.4	0.7
Accidents involving falls	–	–	–	–	–	–	–	–
Accidental drowning or ingestion causing obstruction to the airways	1	1	3	4	0.1	0.1	0.4	0.5
Accidents involving suffocation or strangulation in bed	75	73	60	54	9.7	9.5	8.0	7.1
Assault and other intentional injuries	13	6	9	13	1.7	0.8	1.2	1.7
Injuries with undetermined intent or unspecified cause	3	1	7	8	0.4	0.1	0.9	1.0
Other unintentional injuries	11	21	16	19	1.4	2.7	2.1	2.5
Neoplasms, anemias, and other conditions not otherwise specified	124	134	127	110	16.0	17.4	17.0	14.4

Note: The categories listed in this table conform to "[Infant Mortality: Development of a Proposed Update to the Dollfus Classification of Infant Deaths](#)" (Nakamura, et. al.), with the exception of certain external causes. Due to the high number of accidental suffocations of infants in bed, the DVRHS lists "Accidents involving suffocation or strangulation in bed" as a separate category.

Source: 2014-2017 Michigan Resident Infant Death File, Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services.

Table 15-A
Sudden Infant Death Syndrome (SIDS), Accidental Suffocation & Strangulation in Bed (ASSB)
and
Sudden Unexpected Infant Deaths (SUID),
Michigan Residents 1989-2017

Year	Number				Rate Per 10,000 Live Births		
	SIDS	ASSB	Unknown & unspecified cause	SUID	SIDS	ASSB	SUID
1989	254	10	1	265	17.1	0.7	17.9
1990	256	11	2	269	16.7	0.7	17.6
1991	240	13	3	256	16.1	0.9	17.1
1992	243	7	4	254	16.9	0.5	17.7
1993	210	10	10	230	15.0	0.7	16.5
1994	167	10	8	185	12.1	0.7	13.4
1995	148	15	5	168	11.0	1.1	12.5
1996	149	8	5	162	11.2	0.6	12.2
1997	138	9	13	160	10.3	0.7	12.0
1998	148	11	13	172	11.1	0.8	12.9
1999	114	12	9	135	8.5	0.9	10.1
2000	99	28	6	133	7.3	2.1	9.8
2001	96	39	1	136	7.2	2.9	10.2
2002	85	32	10	127	6.6	2.5	9.8
2003	49	33	14	96	3.7	2.5	7.3
2004	51	49	10	110	3.9	3.8	8.5
2005	47	52	10	109	3.7	4.1	8.5
2006	51	52	9	112	4.0	4.1	8.8
2007	46	56	8	110	3.7	4.5	8.8
2008	48	52	2	102	4.0	4.3	8.4
2009	45	63	4	112	3.8	5.4	9.5
2010	44	52	7	103	3.8	4.5	9.0
2011	33	69	13	115	2.9	6.0	10.1
2012	15	75	16	106	1.3	6.7	9.4
2013	31	72	14	117	2.7	6.3	10.3
2014	35	76	14	125	3.1	6.6	10.9
2015	33	73	24	125	2.9	6.5	11.0
2016	45	60	13	118	4.0	5.3	10.4
2017	32	54	17	99	2.9	4.8	8.9

Table 15-B
Accidental Suffocation & Strangulation in Bed and Total Accidents,
Michigan Residents, 1989-2017

Year	ASSB	Other Accidents & adverse effects	Total Accidents	ASSB Percent of Accidents
1989	10	37	47	21.3%
1990	11	27	38	28.9%
1991	13	25	38	34.2%
1992	7	22	29	24.1%
1993	10	35	45	22.2%
1994	10	27	37	27.0%
1995	15	25	40	37.5%
1996	8	28	36	22.2%
1997	9	24	33	27.3%
1998	11	24	35	31.4%
1999	12	24	36	33.3%
2000	28	27	55	50.9%
2001	39	18	57	68.4%
2002	32	40	72	44.4%
2003	33	31	64	51.6%
2004	49	21	70	70.0%
2005	52	16	68	76.5%
2006	52	19	71	73.2%
2007	56	21	77	72.7%
2008	52	10	62	83.9%
2009	63	11	74	85.1%
2010	52	13	65	80.0%
2011	69	14	83	83.1%
2012	75	17	92	81.5%
2013	72	18	90	80.0%
2014	76	14	90	84.4%
2015	73	24	97	75.3%
2016	60	22	82	73.2%
2017	54	28	82	65.9%

Notes: Causes of death are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. Before 1999, causes of death were classified according to the following ICD-9 codes: SIDS: 798.0, ASSB: E913.0, Unknown: 799.9. On or after 1999, deaths were classified according to the following ICD-10 codes: SIDS: R95, ASSB: W75 and Unknown: R99.

After 1996, SUID death-scene guidelines and subsequent reporting instructions have shifted how SIDS and ASSB deaths were coded. As a result, some deaths that may have been attributed to SIDS before 1999 were classified as ASSB after 1999.

For further information on sudden unexplained infant deaths, see the Sudden Unexpected Infant Death Case Registry. (www.cdc.gov/sids/index.htm)

Table 16
Three-year Average Mortality Rates by Race
Sudden Infant Death Syndrome (SIDS), Accidental Suffocation & Strangulation in Bed (ASSB)
and Sudden Unexpected Infant Deaths (SUID),
Michigan Residents 1989-2017

Trend	All			White			Black		
	SIDS	ASSB	SUIDS	SIDS	ASSB	SUIDS	SIDS	ASSB	SUIDS
1989-1991	16.6	0.8	17.5	13.4	0.5	14.1	27.1	1.7	29.0
1990-1992	16.6	0.7	17.5	13.3	0.6	14.1	28.1	1.1	29.3
1991-1993	16.0	0.7	17.1	12.8	0.7	13.8	28.2	0.8	29.5
1992-1994	14.7	0.6	15.9	11.6	0.6	12.7	27.3	0.8	28.7
1993-1995	12.8	0.9	14.2	9.7	0.6	10.9	25.9	1.7	28.3
1994-1996	11.4	0.8	12.7	8.2	0.6	9.3	26.1	1.6	28.1
1995-1997	10.8	0.8	12.2	7.7	0.7	9.0	24.9	1.2	26.6
1996-1998	10.9	0.7	12.3	7.7	0.6	9.0	24.3	1.1	26.4
1997-1999	10.0	0.8	11.7	7.3	0.6	8.5	21.7	1.7	24.9
1998-2000	9.0	1.3	10.9	6.4	0.9	7.7	19.7	3.1	24.1
1999-2001	7.7	2.0	10.0	5.3	1.6	7.2	18.3	3.8	22.8
2000-2002	7.0	2.5	9.9	5.0	1.9	7.3	16.3	5.6	22.5
2001-2003	5.8	2.6	9.1	4.5	2.2	7.2	12.8	5.3	19.4
2002-2004	4.7	2.9	8.5	3.9	2.2	6.9	9.2	7.0	17.9
2003-2005	3.8	3.5	8.1	3.1	2.9	6.7	7.3	6.7	15.6
2004-2006	3.9	4.0	8.6	3.0	3.0	6.7	7.7	8.9	18.0
2005-2007	3.8	4.2	8.7	3.2	3.1	6.7	6.4	10.1	18.2
2006-2008	3.9	4.3	8.7	3.5	2.9	6.7	5.3	10.8	17.4
2007-2009	3.8	4.7	8.9	3.6	3.0	6.8	4.9	11.9	18.1
2008-2010	3.9	4.7	9.0	3.6	3.1	7.0	4.9	11.0	17.0
2009-2011	3.5	5.3	9.5	3.2	3.5	7.1	5.1	12.8	19.5
2010-2012	2.7	5.7	9.5	2.5	3.9	7.3	3.7	13.7	18.6
2011-2013	2.3	6.3	9.9	2.2	4.3	7.6	2.9	15.7	20.6
2012-2014	2.4	5.8	9.4	2.4	3.7	6.9	2.5	15.1	19.2
2013-2015	2.9	5.7	10.1	2.9	4.0	8.0	3.4	13.3	20.2
2014-2016	3.3	6.1	10.9	3.0	4.5	8.5	4.7	13.8	22.6
2015-2017	3.3	5.5	10.3	2.7	3.7	7.4	5.9	13.3	22.7

Notes: Rates are per 10,000 live births. Causes of death are coded using the 9th and 10th revisions to the International Classification of Diseases. The ICD-9 and ICD-10 codes are grouped into broader categories for the causes listed in this table in order to classify these selected causes of death. Before 1999, causes of death were classified according to the following ICD-9 codes: SIDS: 798.0, ASSB: E913.0, Unknown: 799.9. On or after 1999, deaths were classified according to the following ICD-10 codes: SIDS: R95, ASSB: W75 and Unknown: R99.

After 1996, SUID death-scene guidelines and subsequent reporting instructions shifted how SIDS and ASSB deaths were coded. As a result, some deaths that may have been attributed to SIDS before 1999 were classified as ASSB after 1999.

Table 17-A
Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2006-2017

County	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	7.6±0.3	7.6±0.3	7.3±0.3	7.1±0.3	6.9±0.3	6.8±0.3	6.9±0.3	6.9±0.3	6.7±0.3	6.7±0.3
Alcona	*	*	*	*	*	*	*	*	*	*
Alger	*	*	*	*	-	-	-	-	-	-
Allegan	6.9±2.4	6.7±2.4	4.2±1.9	4.5±2.0	4.2±2.0	5.0±2.2	5.4±2.3	6.8±2.5	5.7±2.3	4.6±2.1
Alpena	6.7±5.3	*	*	7.3±5.8	*	*	*	*	*	*
Antrim	8.7±6.9	*	11.1±8.2	10.9±8.0	10.0±8.0	*	*	*	-	-
Arenac	*	*	*	*	*	16.2±12.9	16.5±13.1	*	*	*
Baraga	*	*	*	-	-	*	*	*	*	*
Barry	7.8±3.7	7.4±3.7	8.9±4.1	7.9±4.0	6.8±3.7	4.9±3.2	4.8±3.1	3.2±2.5	*	3.6±2.7
Bay	7.5±2.8	6.6±2.6	7.9±2.9	8.3±3.0	7.7±3.0	5.3±2.5	5.7±2.6	6.1±2.7	5.4±2.6	4.5±2.4
Benzie	*	*	*	*	*	*	*	*	*	*
Berrien	8.8±2.3	8.9±2.4	7.8±2.2	8.3±2.3	8.1±2.3	8.3±2.4	7.8±2.3	6.9±2.2	6.4±2.2	5.2±2.0
Branch	3.3±2.6	3.5±2.8	4.9±3.4	6.3±3.9	7.8±4.4	7.0±4.1	6.9±4.1	6.8±4.0	8.1±4.4	8.7±4.6
Calhoun	10.8±2.7	11.5±2.9	10.2±2.8	10.7±2.9	8.7±2.6	7.2±2.3	6.0±2.1	6.8±2.3	5.8±2.1	7.2±2.4
Cass	5.9±3.6	7.8±4.2	7.8±4.4	7.3±4.3	6.6±4.3	7.4±4.6	8.5±4.8	7.3±4.3	7.3±4.3	6.2±4.0
Charlevoix	*	*	*	*	*	*	*	*	*	*
Cheboygan	*	*	*	*	-	-	-	*	*	*
Chippewa	5.4±4.3	*	*	5.3±4.2	7.5±5.2	7.5±5.2	5.6±4.5	*	*	*
Clare	6.6±4.9	5.8±4.7	*	*	*	*	*	*	*	6.1±4.9
Clinton	4.8±2.7	4.7±2.8	4.4±2.7	4.0±2.6	3.8±2.5	2.5±2.0	3.8±2.5	6.3±3.2	6.5±3.2	5.2±2.8
Crawford	19.1±13.1	17.5±12.9	*	*	*	*	*	*	*	15.7±12.5
Delta	*	6.1±4.5	*	5.3±4.2	6.4±4.7	5.4±4.3	7.1±4.9	*	5.4±4.3	*
Dickinson	7.6±6.0	*	8.3±6.6	*	*	*	*	*	*	*
Eaton	5.3±2.4	5.3±2.4	5.9±2.5	5.6±2.5	5.4±2.4	6.1±2.6	6.0±2.6	7.0±2.8	6.7±2.7	6.0±2.5
Emmet	*	*	6.3±5.0	*	*	*	*	*	*	*
Genesee	9.3±1.4	9.0±1.4	7.8±1.4	7.8±1.4	7.1±1.3	8.0±1.4	8.0±1.4	8.5±1.5	9.1±1.5	9.0±1.5
Gladwin	*	*	*	*	*	*	*	11.2±7.3	9.7±6.7	*
Gogebic	*	*	*	*	*	-	*	*	16.1±12.8	*
Grand Traverse	7.0±3.0	6.9±3.0	5.6±2.8	3.3±2.1	2.9±2.0	*	3.5±2.2	5.3±2.7	7.5±3.2	6.5±3.0
Gratiot	8.7±4.9	7.1±4.4	8.1±4.7	8.1±4.8	6.3±4.3	4.8±3.8	5.8±4.2	6.6±4.6	5.1±4.0	*
Hillsdale	9.2±4.5	5.9±3.6	4.2±3.1	*	3.7±3.0	4.3±3.2	*	4.8±3.3	7.3±4.1	11.2±5.1

Table 17-A
Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2006-2017

County	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Houghton	7.3±4.8	6.6±4.5	6.5±4.5	*	5.1±4.0	5.3±4.3	5.4±4.3	*	*	6.3±4.7
Huron	*	*	*	6.3±5.1	6.5±5.2	*	*	-	*	*
Ingham	6.6±1.6	7.0±1.6	5.9±1.5	5.6±1.5	5.8±1.5	7.2±1.7	7.1±1.7	7.1±1.7	6.4±1.6	6.0±1.5
Ionia	3.9±2.4	3.3±2.3	3.0±2.2	4.0±2.6	4.0±2.6	4.1±2.6	4.1±2.7	5.0±2.9	4.6±2.8	4.1±2.7
Iosco	*	*	11.7±8.6	12.6±8.6	9.2±7.4	*	*	*	*	8.8±7.0
Iron	*	*	*	*	*	*	*	*	*	*
Isabella	6.2±3.2	5.4±3.1	4.2±2.7	6.8±3.5	7.1±3.7	6.2±3.5	4.0±2.8	4.4±2.9	4.0±2.7	7.0±3.7
Jackson	7.7±2.2	7.1±2.2	6.5±2.1	6.1±2.1	5.3±1.9	6.2±2.1	5.6±2.0	6.2±2.1	6.5±2.1	7.8±2.4
Kalamazoo	7.1±1.7	8.4±1.8	8.6±1.9	7.6±1.8	6.4±1.6	6.0±1.6	6.4±1.6	6.5±1.6	4.6±1.4	6.7±1.6
Kalkaska	-	*	*	*	11.3±9.0	11.6±9.2	17.0±11.0	*	*	*
Kent	7.8±1.0	6.9±1.0	6.8±1.0	6.4±1.0	6.2±0.9	5.6±0.9	5.3±0.9	5.5±0.9	6.2±0.9	6.0±0.9
Keweenaw	*	-	-	-	-	*	*	*	-	-
Lake	*	*	*	*	*	*	*	*	*	*
Lapeer	5.4±2.7	7.4±3.2	8.6±3.6	7.1±3.3	3.6±2.4	2.4±1.9	4.4±2.6	5.3±2.9	5.3±2.9	4.6±2.7
Leelanau	*	*	*	*	*	*	*	*	*	*
Lenawee	8.3±3.0	8.5±3.1	10.8±3.5	9.4±3.3	9.2±3.3	7.2±2.9	8.2±3.1	7.6±3.0	7.3±3.0	6.2±2.8
Livingston	5.5±1.9	6.7±2.2	6.3±2.1	6.7±2.2	5.9±2.1	5.2±2.0	5.4±2.0	5.1±1.9	4.4±1.7	3.5±1.6
Luce	*	*	*	-	-	-	*	*	*	*
Mackinac	*	*	*	*	-	-	-	*	*	*
Macomb	6.1±0.9	7.3±1.0	6.7±1.0	6.5±1.0	5.6±0.9	6.3±0.9	6.0±0.9	6.3±0.9	5.7±0.9	6.4±0.9
Manistee	*	*	*	*	*	*	*	*	*	*
Marquette	4.0±2.8	3.9±2.7	4.1±2.8	5.1±3.1	4.1±2.8	4.1±2.9	*	4.3±3.0	4.2±2.9	4.3±3.0
Mason	7.0±5.2	*	6.4±5.1	*	*	*	*	*	*	*
Mecosta	9.7±5.2	8.2±4.8	9.0±5.1	7.0±4.5	10.2±5.5	11.8±6.0	9.3±5.2	8.6±5.1	5.5±4.1	*
Menominee	*	*	*	*	*	*	*	*	*	*
Midland	5.3±2.8	4.2±2.5	3.9±2.4	4.2±2.5	4.6±2.6	6.6±3.1	7.1±3.3	5.8±2.9	3.5±2.3	2.3±1.9
Missaukee	-	*	*	*	*	*	*	*	*	*
Monroe	6.7±2.2	5.7±2.0	5.7±2.1	4.3±1.8	4.4±1.9	5.8±2.2	5.9±2.2	6.0±2.3	4.2±1.9	6.1±2.3
Montcalm	3.7±2.4	3.9±2.6	5.8±3.1	6.4±3.3	6.8±3.4	6.4±3.3	5.5±3.1	4.6±2.9	4.3±2.8	4.7±2.9
Montmorency	*	-	*	*	*	*	*	-	-	-

Table 17-A
Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2006-2017

County	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Muskegon	6.7±1.9	7.1±2.0	7.9±2.1	8.0±2.1	8.7±2.3	8.0±2.2	8.1±2.2	8.1±2.2	8.4±2.3	8.4±2.2
Newaygo	8.4±4.1	8.2±4.1	5.3±3.5	7.8±4.2	7.3±4.1	7.3±4.1	7.2±4.1	7.9±4.3	7.9±4.3	6.2±3.8
Oakland	6.4±0.8	6.2±0.8	6.2±0.8	5.8±0.7	6.2±0.8	6.3±0.8	6.6±0.8	6.0±0.8	5.6±0.7	5.3±0.7
Oceana	6.0±4.5	6.3±4.6	8.3±5.4	5.9±4.7	9.0±5.9	*	12.3±7.2	9.1±6.2	12.1±7.1	*
Ogemaw	*	*	*	*	*	*	*	*	*	*
Ontonagon	*	*	-	-	-	-	*	*	*	-
Osceola	*	*	*	*	*	*	*	*	*	8.0±6.4
Oscoda	*	*	*	*	*	*	*	*	*	-
Otsego	9.3±6.4	8.6±6.3	*	9.0±6.6	11.9±7.7	13.9±8.6	12.6±8.2	9.3±6.9	*	*
Ottawa	5.9±1.5	5.4±1.4	5.7±1.5	5.6±1.5	5.9±1.5	6.6±1.6	5.7±1.5	6.0±1.5	4.9±1.4	4.8±1.4
Presque Isle	*	*	*	*	21.7±17.1	*	*	*	*	*
Roscommon	*	10.9±8.7	*	*	*	*	*	*	*	*
Saginaw	9.4±2.2	10.5±2.3	8.8±2.2	7.9±2.1	6.1±1.8	7.1±2.0	7.7±2.1	8.7±2.2	10.0±2.4	8.4±2.2
St. Clair	5.8±2.0	6.4±2.1	5.6±2.0	6.3±2.2	6.2±2.2	6.3±2.2	6.0±2.2	5.1±2.1	6.2±2.2	5.9±2.2
St. Joseph	10.3±3.7	11.2±4.0	10.6±4.0	8.3±3.5	7.4±3.4	7.3±3.4	7.9±3.5	5.3±2.8	5.1±2.9	4.8±2.8
Sanilac	7.5±4.4	6.8±4.2	*	*	*	4.6±3.6	6.8±4.4	6.8±4.4	8.4±4.9	8.4±4.9
Schoolcraft	*	*	-	-	-	-	-	-	-	-
Shiawassee	5.4±2.9	5.7±3.1	3.2±2.4	3.3±2.4	3.8±2.6	3.2±2.4	4.2±2.7	4.7±2.9	6.1±3.3	7.2±3.6
Tuscola	3.8±2.8	3.9±2.9	6.0±3.7	6.8±4.0	6.8±4.0	4.9±3.4	4.7±3.3	4.7±3.3	4.2±3.1	4.2±3.1
Van Buren	5.5±2.6	6.9±2.9	6.5±2.9	4.5±2.4	5.6±2.8	6.5±3.0	6.2±3.0	5.6±2.8	6.1±3.0	6.2±3.0
Washtenaw	5.2±1.3	5.3±1.3	5.1±1.3	4.9±1.3	4.7±1.3	4.6±1.3	5.9±1.4	5.5±1.4	5.9±1.4	4.7±1.3
Wayne	10.4±0.7	10.4±0.7	10.2±0.7	9.8±0.7	9.9±0.7	9.7±0.7	9.6±0.7	9.4±0.7	9.3±0.7	9.7±0.7
Wexford	6.8±4.4	7.7±4.8	6.3±4.3	6.4±4.4	9.8±5.5	9.8±5.5	10.8±5.8	6.3±4.4	5.0±4.0	5.1±4.1

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

**Table 17-B
White Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2006-2017**

County	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	5.6±0.3	5.6±0.3	5.4±0.3	5.2±0.3	5.3±0.3	5.3±0.3	5.5±0.3	5.4±0.3	5.1±0.3	4.9±0.3
Alcona	*	*	*	*	*	*	*	*	*	-
Alger	*	*	*	*	-	-	-	-	-	-
Allegan	5.9±2.3	5.6±2.2	3.4±1.8	4.0±2.0	4.1±2.0	5.0±2.2	5.4±2.3	6.8±2.6	5.4±2.3	4.5±2.1
Alpena	6.9±5.5	*	*	*	*	*	*	*	*	*
Antrim	8.9±7.1	*	11.7±8.6	11.7±8.6	10.7±8.5	*	*	*	-	-
Arenac	*	*	*	*	*	16.6±13.2	16.8±13.3	*	*	*
Baraga	-	-	-	-	-	-	*	*	*	*
Barry	8.0±3.8	7.5±3.8	9.1±4.2	8.0±4.0	6.9±3.7	5.0±3.3	4.9±3.2	3.2±2.6	*	3.7±2.7
Bay	7.1±2.8	6.1±2.6	7.4±2.9	7.9±3.0	7.6±3.0	5.3±2.6	5.5±2.7	5.9±2.8	5.1±2.6	4.8±2.5
Benzie	*	*	*	*	*	*	*	*	*	*
Berrien	5.9±2.3	4.8±2.1	4.0±1.9	3.6±1.8	4.6±2.1	4.9±2.2	4.9±2.2	4.7±2.1	5.1±2.3	4.4±2.2
Branch	3.4±2.7	3.6±2.9	5.0±3.5	6.5±4.0	6.7±4.1	5.2±3.6	5.2±3.6	6.5±4.0	8.6±4.6	7.8±4.4
Calhoun	7.9±2.6	9.3±2.9	7.4±2.6	7.9±2.8	6.4±2.5	5.9±2.4	5.1±2.2	5.2±2.3	4.4±2.1	6.2±2.5
Cass	5.5±3.8	5.6±3.9	5.9±4.1	5.3±3.9	6.8±4.7	7.6±5.0	9.6±5.4	8.1±4.8	8.1±4.8	6.1±4.2
Charlevoix	*	*	*	*	*	*	*	*	*	*
Cheboygan	*	*	*	*	-	-	-	*	*	*
Chippewa	*	*	*	*	*	*	*	*	*	*
Clare	6.8±5.0	6.0±4.8	*	*	*	*	*	*	*	6.3±5.0
Clinton	4.7±2.8	4.5±2.8	4.6±2.8	4.2±2.7	4.1±2.7	2.7±2.2	4.1±2.7	5.9±3.2	6.3±3.3	4.4±2.7
Crawford	17.0±12.5	15.7±12.5	*	*	*	*	-	*	*	16.4±13.0
Delta	*	*	*	*	5.8±4.7	*	6.5±4.8	*	*	*
Dickinson	8.0±6.4	*	*	*	*	*	*	*	*	*
Eaton	5.1±2.5	3.8±2.1	4.2±2.3	3.9±2.2	4.8±2.5	4.8±2.5	4.8±2.5	5.1±2.6	5.1±2.6	4.5±2.5
Emmet	*	*	6.8±5.4	*	*	*	*	*	*	*
Genesee	6.0±1.4	5.4±1.3	5.1±1.3	5.1±1.4	4.8±1.3	5.2±1.4	5.3±1.4	5.7±1.5	6.3±1.6	6.4±1.6
Gladwin	*	*	*	*	*	*	*	11.4±7.4	9.9±6.8	*
Gogebic	*	*	*	*	*	-	-	-	*	*
Grand Traverse	7.4±3.2	7.3±3.2	6.0±3.0	3.5±2.3	3.1±2.2	*	3.7±2.3	5.6±2.8	7.8±3.3	6.8±3.1
Gratiot	7.7±4.8	6.0±4.2	7.0±4.5	7.8±4.8	5.8±4.3	*	5.2±4.2	6.2±4.6	5.4±4.3	*
Hillsdale	9.6±4.7	6.2±3.8	4.3±3.2	*	3.8±3.1	4.4±3.2	*	4.9±3.4	7.5±4.2	11.4±5.2

Table 17-B
White Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2006-2017

County	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Houghton	7.0±4.8	6.1±4.5	6.0±4.4	*	5.4±4.3	5.7±4.6	5.7±4.6	*	*	5.8±4.6
Huron	*	*	*	6.5±5.2	6.7±5.3	*	*	-	*	*
Ingham	4.1±1.4	4.7±1.5	3.6±1.4	4.6±1.6	5.3±1.7	7.5±2.1	7.1±2.0	6.4±1.9	4.6±1.6	4.3±1.6
Ionia	4.1±2.5	3.4±2.4	3.1±2.3	3.6±2.5	3.7±2.6	3.7±2.6	4.2±2.8	5.1±3.0	4.8±2.9	4.3±2.8
Iosco	*	*	12.2±9.0	13.1±9.0	9.6±7.6	*	*	*	*	*
Iron	*	*	*	*	*	*	*	*	*	*
Isabella	5.0±3.1	3.1±2.5	*	5.6±3.4	6.5±3.8	5.9±3.7	3.5±2.8	4.0±3.0	4.0±3.0	7.1±4.0
Jackson	6.4±2.2	6.2±2.2	5.7±2.1	5.5±2.1	4.1±1.8	4.6±2.0	3.9±1.8	5.1±2.1	4.8±2.0	5.6±2.2
Kalamazoo	4.8±1.6	5.5±1.7	5.7±1.7	4.8±1.6	3.9±1.4	3.9±1.4	4.7±1.6	4.7±1.6	3.2±1.3	4.7±1.6
Kalkaska	-	*	*	*	11.7±9.3	11.9±9.5	17.4±11.3	*	*	*
Kent	5.6±1.0	5.0±0.9	5.0±0.9	4.4±0.9	4.4±0.9	4.1±0.9	4.5±0.9	4.9±0.9	5.1±1.0	4.8±0.9
Keweenaw	*	-	-	-	-	*	*	*	-	-
Lake	*	*	*	*	*	*	*	*	*	*
Lapeer	5.5±2.8	7.3±3.3	7.8±3.5	6.3±3.2	3.0±2.2	2.6±2.1	4.7±2.8	5.6±3.0	5.6±3.0	4.8±2.9
Leelanau	*	*	*	*	*	*	*	*	*	*
Lenawee	9.0±3.3	8.4±3.3	10.6±3.7	8.7±3.4	9.6±3.6	7.6±3.2	8.4±3.3	7.6±3.2	7.0±3.1	6.5±3.0
Livingston	5.3±1.9	6.6±2.2	6.1±2.2	6.6±2.2	5.8±2.1	5.0±2.0	5.1±2.0	4.7±1.9	4.2±1.7	3.0±1.5
Luce	*	*	*	-	-	-	*	*	*	*
Mackinac	*	*	*	*	-	-	-	*	*	*
Macomb	5.2±0.9	6.2±1.0	5.9±1.0	5.6±1.0	4.8±0.9	5.5±1.0	5.3±1.0	5.4±1.0	4.6±0.9	5.2±1.0
Manistee	*	*	*	*	*	*	*	*	*	*
Marquette	4.2±2.9	3.6±2.7	3.7±2.8	4.8±3.1	4.2±2.9	4.3±3.0	*	4.5±3.1	4.4±3.0	4.5±3.1
Mason	6.2±4.9	*	6.6±5.3	*	*	*	*	*	*	*
Mecosta	9.4±5.3	7.9±4.9	8.8±5.2	6.6±4.5	10.0±5.6	11.7±6.1	9.9±5.6	9.1±5.4	5.8±4.3	*
Menominee	*	*	*	*	*	*	*	*	*	-
Midland	5.6±2.9	4.0±2.5	3.7±2.4	4.0±2.5	4.9±2.8	5.8±3.0	6.3±3.2	4.9±2.8	3.7±2.4	2.5±2.0
Missaukee	-	*	*	*	*	*	*	*	*	*
Monroe	6.4±2.3	4.8±2.0	4.7±1.9	3.5±1.7	3.6±1.8	5.1±2.1	5.3±2.2	6.0±2.3	4.1±1.9	6.1±2.4
Montcalm	3.8±2.5	4.0±2.6	5.9±3.2	6.1±3.3	6.5±3.4	6.0±3.3	5.6±3.2	4.7±2.9	4.4±2.8	4.8±3.0
Montmorency	*	-	*	*	*	*	*	-	-	-

**Table 17-B
White Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2006-2017**

County	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Muskegon	5.7±2.0	6.4±2.2	6.5±2.2	6.3±2.2	7.1±2.4	6.6±2.3	5.8±2.2	4.6±1.9	4.1±1.8	4.2±1.8
Newaygo	8.0±4.0	8.4±4.2	5.4±3.5	7.4±4.2	6.9±4.1	7.0±4.1	7.6±4.3	8.4±4.6	8.4±4.5	6.6±4.1
Oakland	4.9±0.8	4.8±0.8	5.3±0.8	5.1±0.8	5.7±0.9	5.6±0.9	5.4±0.8	4.5±0.8	4.0±0.7	3.8±0.7
Oceana	*	*	5.9±4.7	*	9.6±6.3	*	13.5±7.9	8.7±6.4	10.7±7.0	*
Ogemaw	*	*	*	*	*	*	*	*	*	*
Ontonagon	*	*	-	-	-	-	*	*	*	-
Osceola	*	*	*	*	*	*	*	*	*	*
Oscoda	*	*	*	*	*	*	*	*	*	-
Otsego	9.4±6.5	8.6±6.4	*	9.2±6.8	12.2±7.9	14.4±8.9	11.5±8.0	8.3±6.6	*	*
Ottawa	5.3±1.4	4.6±1.4	4.9±1.4	5.3±1.5	5.6±1.5	6.3±1.6	5.3±1.5	5.6±1.5	4.7±1.4	4.6±1.4
Presque Isle	*	*	*	*	22.1±17.5	*	*	*	*	*
Roscommon	*	11.2±8.9	*	*	*	*	*	*	*	*
Saginaw	6.6±2.2	6.4±2.2	4.7±1.9	2.9±1.5	2.9±1.5	4.2±1.8	6.2±2.2	6.3±2.2	7.5±2.5	5.5±2.1
St. Clair	5.0±1.9	6.2±2.1	5.4±2.1	6.3±2.3	6.2±2.3	6.3±2.3	5.9±2.3	4.3±1.9	5.0±2.1	4.7±2.1
St. Joseph	9.3±3.6	10.2±3.9	9.9±3.9	7.9±3.6	6.9±3.4	6.4±3.2	6.6±3.3	4.3±2.7	4.5±2.8	4.7±2.9
Sanilac	7.0±4.3	6.3±4.1	*	*	*	4.6±3.7	6.9±4.5	6.1±4.2	7.7±4.8	7.8±4.8
Schoolcraft	*	*	-	-	-	-	-	-	-	-
Shiawassee	5.5±3.0	5.9±3.2	3.3±2.4	3.4±2.5	3.9±2.7	3.3±2.5	4.3±2.8	4.3±2.8	5.8±3.3	6.9±3.6
Tuscola	3.9±2.9	4.0±3.0	6.1±3.8	6.9±4.1	7.0±4.1	5.0±3.5	4.9±3.4	4.8±3.3	4.3±3.2	4.4±3.2
Van Buren	4.8±2.6	6.1±3.0	6.0±3.0	4.1±2.5	5.4±2.9	6.2±3.1	5.8±3.0	5.3±2.9	6.2±3.1	6.3±3.2
Washtenaw	3.5±1.3	4.3±1.4	4.4±1.5	4.5±1.5	3.9±1.4	4.2±1.5	5.4±1.7	5.0±1.6	4.3±1.5	3.2±1.3
Wayne	5.6±0.7	5.6±0.8	5.4±0.8	5.5±0.8	5.9±0.9	6.3±0.9	6.4±0.9	6.4±0.9	6.1±0.9	5.6±0.8
Wexford	7.0±4.6	8.0±5.0	6.5±4.5	6.6±4.5	10.1±5.7	9.3±5.4	10.3±5.8	5.7±4.2	5.1±4.1	5.3±4.2

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 18-C
Black Three-Year Moving Average Infant Death Rates by County
Michigan Residents, 2006-2017

County	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	15.3±0.9	15.5±0.9	14.8±0.9	14.5±0.9	13.8±0.9	13.4±0.9	13.3±0.9	13.7±0.9	13.8±0.9	14.3±0.9
Berrien	14.9±6.0	16.7±6.4	16.0±6.2	19.5±6.9	18.6±6.9	17.3±6.7	16.5±6.7	13.3±6.1	10.4±5.6	7.4±4.8
Calhoun	24.2±10.5	20.5±9.6	22.1±10.4	25.4±11.3	20.2±10.1	13.8±8.1	9.9±6.8	14.6±8.2	11.3±7.3	12.6±7.8
Cass	*	*	*	*	*	*	-	-	-	-
Genesee	17.1±3.6	17.9±3.7	14.6±3.4	14.2±3.4	12.4±3.3	14.2±3.5	14.6±3.6	15.6±3.7	16.5±3.9	15.2±3.7
Ingham	17.8±6.3	16.5±6.0	16.7±5.9	13.1±5.3	11.4±5.0	10.1±4.7	10.1±4.7	12.9±5.2	14.2±5.5	14.5±5.6
Jackson	16.8±10.3	15.6±10.1	13.2±9.1	11.6±8.6	13.0±9.0	18.5±10.8	16.2±10.0	15.0±9.7	20.2±11.3	26.5±12.8
Kalamazoo	16.8±6.2	19.2±6.5	19.5±6.5	18.3±6.4	17.6±6.3	15.5±5.9	14.2±5.7	15.3±5.9	11.0±4.9	15.3±5.6
Kent	14.4±3.8	13.4±3.7	13.9±3.8	16.6±4.2	14.6±3.9	12.9±3.6	9.9±3.2	10.7±3.3	13.0±3.7	12.8±3.6
Macomb	13.4±4.0	15.1±4.0	13.8±3.8	13.1±3.6	11.4±3.2	11.1±3.1	9.9±2.9	10.7±2.9	10.6±2.9	12.8±3.1
Muskegon	9.0±4.9	8.2±4.6	13.2±5.9	14.9±6.3	16.4±6.8	12.8±6.0	14.9±6.7	19.7±7.7	23.1±8.3	23.2±8.1
Oakland	15.4±3.1	14.7±3.0	11.9±2.7	10.8±2.5	10.6±2.4	11.1±2.5	13.7±2.8	14.7±2.9	14.3±2.9	13.5±2.8
Saginaw	15.3±5.2	19.2±5.8	17.4±5.6	19.2±6.1	13.4±5.1	14.3±5.2	12.2±5.0	15.9±5.7	17.3±5.9	16.2±5.6
St. Clair	*	*	*	*	*	*	*	*	34.3±25.0	33.7±24.5
Van Buren	*	*	*	*	*	*	*	*	*	*
Washtenaw	13.2±4.9	11.6±4.6	10.5±4.4	8.7±4.0	8.9±4.1	8.3±4.1	10.9±4.7	12.1±5.2	16.4±5.9	14.4±5.5
Wayne	15.3±1.3	15.3±1.3	15.1±1.3	14.5±1.3	14.5±1.3	14.0±1.3	13.7±1.3	13.2±1.2	13.2±1.3	14.1±1.3

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 18-A
Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2006-2017

Health Department	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	7.6±0.3	7.6±0.3	7.3±0.3	7.1±0.3	6.9±0.3	6.8±0.3	6.9±0.3	6.9±0.3	6.7±0.3	6.7±0.3
Allegan County	6.9±2.4	6.7±2.4	4.2±1.9	4.5±2.0	4.2±2.0	5.0±2.2	5.4±2.3	6.8±2.5	5.7±2.3	4.6±2.1
Barry-Eaton District	6.2±2.0	6.0±2.0	7.0±2.2	6.4±2.1	5.9±2.0	5.7±2.0	5.6±2.0	5.7±2.0	5.0±1.9	5.1±1.9
Bay County	7.5±2.8	6.6±2.6	7.9±2.9	8.3±3.0	7.7±3.0	5.3±2.5	5.7±2.6	6.1±2.7	5.4±2.6	4.5±2.4
Benzie-Leelanau District	*	*	*	*	*	*	6.2±5.0	8.2±5.7	8.2±5.7	6.0±4.8
Berrien County	8.8±2.3	8.9±2.4	7.8±2.2	8.3±2.3	8.1±2.3	8.3±2.4	7.8±2.3	6.9±2.2	6.4±2.2	5.2±2.0
Branch-Hillsdale-St. Joseph	8.0±2.2	7.5±2.2	7.2±2.2	6.3±2.0	6.4±2.1	6.3±2.1	6.2±2.0	5.6±1.9	6.6±2.1	7.8±2.3
Calhoun County	10.8±2.7	11.5±2.9	10.2±2.8	10.7±2.9	8.7±2.6	7.2±2.3	6.0±2.1	6.8±2.3	5.8±2.1	7.2±2.4
Central Michigan District	5.6±1.9	5.7±1.9	4.6±1.8	6.3±2.1	5.6±2.0	5.4±2.0	4.6±1.8	5.7±2.0	5.5±2.0	6.3±2.1
Chippewa County	5.4±4.3	*	*	5.3±4.2	7.5±5.2	7.5±5.2	5.6±4.5	*	*	*
City Of Detroit	14.3±1.2	14.8±1.3	14.4±1.3	13.6±1.3	13.6±1.3	13.6±1.3	13.8±1.3	13.6±1.3	13.6±1.3	14.0±1.4
Dickinson-Iron District	6.5±4.8	5.6±4.5	8.7±5.7	8.0±5.5	*	*	5.8±4.6	5.9±4.7	*	*
District #10	6.8±1.7	6.7±1.7	6.7±1.7	6.6±1.7	7.6±1.9	7.0±1.8	8.2±1.9	7.1±1.8	7.0±1.8	5.3±1.6
District #2	5.9±3.6	8.4±4.4	9.8±4.8	9.1±4.6	8.9±4.5	6.5±3.8	7.0±3.9	4.1±3.1	4.6±3.2	5.3±3.4
District #4	6.3±3.3	6.1±3.3	6.1±3.4	6.8±3.7	6.5±3.7	4.9±3.2	4.8±3.1	4.1±2.9	4.6±3.0	6.1±3.4
Genesee County	9.3±1.4	9.0±1.4	7.8±1.4	7.8±1.4	7.1±1.3	8.0±1.4	8.0±1.4	8.5±1.5	9.1±1.5	9.0±1.5
Grand Traverse County	7.0±3.0	6.9±3.0	5.6±2.8	3.3±2.1	2.9±2.0	*	3.5±2.2	5.3±2.7	7.5±3.2	6.5±3.0
Huron County	*	*	*	6.3±5.1	6.5±5.2	*	*	-	*	*
Ingham County	6.6±1.6	7.0±1.6	5.9±1.5	5.6±1.5	5.8±1.5	7.2±1.7	7.1±1.7	7.1±1.7	6.4±1.6	6.0±1.5
Ionia County	3.9±2.4	3.3±2.3	3.0±2.2	4.0±2.6	4.0±2.6	4.1±2.6	4.1±2.7	5.0±2.9	4.6±2.8	4.1±2.7
Jackson County	7.7±2.2	7.1±2.2	6.5±2.1	6.1±2.1	5.3±1.9	6.2±2.1	5.6±2.0	6.2±2.1	6.5±2.1	7.8±2.4
Kalamazoo County	7.1±1.7	8.4±1.8	8.6±1.9	7.6±1.8	6.4±1.6	6.0±1.6	6.4±1.6	6.5±1.6	4.6±1.4	6.7±1.6
Kent County	7.8±1.0	6.9±1.0	6.8±1.0	6.4±1.0	6.2±0.9	5.6±0.9	5.3±0.9	5.5±0.9	6.2±0.9	6.0±0.9
Lapeer County	5.4±2.7	7.4±3.2	8.6±3.6	7.1±3.3	3.6±2.4	2.4±1.9	4.4±2.6	5.3±2.9	5.3±2.9	4.6±2.7
Lenawee County	8.3±3.0	8.5±3.1	10.8±3.5	9.4±3.3	9.2±3.3	7.2±2.9	8.2±3.1	7.6±3.0	7.3±3.0	6.2±2.8
Livingston County	5.5±1.9	6.7±2.2	6.3±2.1	6.7±2.2	5.9±2.1	5.2±2.0	5.4±2.0	5.1±1.9	4.4±1.7	3.5±1.6
Luce-Mackinac-Alger-Schoolcraft	13.2±7.4	11.2±6.9	10.4±6.8	*	-	-	*	*	11.6±7.6	9.0±6.6
Macomb County	6.1±0.9	7.3±1.0	6.7±1.0	6.5±1.0	5.6±0.9	6.3±0.9	6.0±0.9	6.3±0.9	5.7±0.9	6.4±0.9
Marquette County	4.0±2.8	3.9±2.7	4.1±2.8	5.1±3.1	4.1±2.8	4.1±2.9	*	4.3±3.0	4.2±2.9	4.3±3.0
Midland County	5.3±2.8	4.2±2.5	3.9±2.4	4.2±2.5	4.6±2.6	6.6±3.1	7.1±3.3	5.8±2.9	3.5±2.3	2.3±1.9

**Table 18-A
Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2006-2017**

Health Department	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Mid-Michigan District	5.3±1.8	5.0±1.8	5.8±1.9	5.9±2.0	5.5±1.9	4.4±1.7	4.8±1.8	5.7±1.9	5.4±1.9	4.6±1.7
Monroe County	6.7±2.2	5.7±2.0	5.7±2.1	4.3±1.8	4.4±1.9	5.8±2.2	5.9±2.2	6.0±2.3	4.2±1.9	6.1±2.3
Muskegon County	6.7±1.9	7.1±2.0	7.9±2.1	8.0±2.1	8.7±2.3	8.0±2.2	8.1±2.2	8.1±2.2	8.4±2.3	8.4±2.2
Northwest Michigan Community Health	6.1±2.6	6.5±2.8	6.4±2.8	6.8±2.9	6.9±2.9	6.7±2.9	6.2±2.8	5.8±2.7	3.7±2.2	3.5±2.1
Oakland County	6.4±0.8	6.2±0.8	6.2±0.8	5.8±0.7	6.2±0.8	6.3±0.8	6.6±0.8	6.0±0.8	5.6±0.7	5.3±0.7
Ottawa County	5.9±1.5	5.4±1.4	5.7±1.5	5.6±1.5	5.9±1.5	6.6±1.6	5.7±1.5	6.0±1.5	4.9±1.4	4.8±1.4
Delta & Menominee	4.3±3.0	6.0±3.5	4.6±3.2	4.6±3.2	5.4±3.5	4.7±3.2	5.7±3.5	4.0±2.9	5.2±3.4	3.7±2.9
Saginaw County	9.4±2.2	10.5±2.3	8.8±2.2	7.9±2.1	6.1±1.8	7.1±2.0	7.7±2.1	8.7±2.2	10.0±2.4	8.4±2.2
Sanilac County	7.5±4.4	6.8±4.2	*	*	*	4.6±3.6	6.8±4.4	6.8±4.4	8.4±4.9	8.4±4.9
Shiawassee County	5.4±2.9	5.7±3.1	3.2±2.4	3.3±2.4	3.8±2.6	3.2±2.4	4.2±2.7	4.7±2.9	6.1±3.3	7.2±3.6
St Clair County	5.8±2.0	6.4±2.1	5.6±2.0	6.3±2.2	6.2±2.2	6.3±2.2	6.0±2.2	5.1±2.1	6.2±2.2	5.9±2.2
Tuscola County	3.8±2.8	3.9±2.9	6.0±3.7	6.8±4.0	6.8±4.0	4.9±3.4	4.7±3.3	4.7±3.3	4.2±3.1	4.2±3.1
Van Buren-Cass County	5.6±2.1	7.2±2.4	6.9±2.4	5.5±2.2	6.0±2.3	6.8±2.5	7.0±2.5	6.2±2.4	6.6±2.5	6.2±2.4
Washtenaw County	5.2±1.3	5.3±1.3	5.1±1.3	4.9±1.3	4.7±1.3	4.6±1.3	5.9±1.4	5.5±1.4	5.9±1.4	4.7±1.3
Wayne County	6.9±0.8	6.7±0.8	6.6±0.8	6.7±0.8	6.9±0.8	6.8±0.8	6.5±0.8	6.3±0.8	6.1±0.8	6.6±0.8
Western Upper Peninsula District	7.5±3.7	5.2±3.1	5.3±3.1	3.5±2.6	4.1±2.8	4.3±2.9	6.0±3.5	5.4±3.4	7.1±3.8	8.1±4.1

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 19-B
White Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2006-2017

Health Department	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	5.6±0.3	5.6±0.3	5.4±0.3	5.2±0.3	5.3±0.3	5.3±0.3	5.5±0.3	5.4±0.3	5.1±0.3	4.9±0.3
Allegan County	5.9±2.3	5.6±2.2	3.4±1.8	4.0±2.0	4.1±2.0	5.0±2.2	5.4±2.3	6.8±2.6	5.4±2.3	4.5±2.1
Barry-Eaton District	6.3±2.1	5.2±2.0	6.1±2.1	5.5±2.1	5.6±2.1	4.9±2.0	4.9±2.0	4.4±1.9	3.9±1.8	4.2±1.8
Bay County	7.1±2.8	6.1±2.6	7.4±2.9	7.9±3.0	7.6±3.0	5.3±2.6	5.5±2.7	5.9±2.8	5.1±2.6	4.8±2.5
Benzie-Leelanau District	*	*	*	*	*	*	*	6.8±5.4	6.7±5.3	*
Berrien County	5.9±2.3	4.8±2.1	4.0±1.9	3.6±1.8	4.6±2.1	4.9±2.2	4.9±2.2	4.7±2.1	5.1±2.3	4.4±2.2
Branch-Hillsdale-St. Joseph	7.7±2.2	7.2±2.2	6.9±2.2	6.2±2.1	6.0±2.1	5.5±2.0	5.2±1.9	5.1±1.9	6.6±2.2	7.6±2.4
Calhoun County	7.9±2.6	9.3±2.9	7.4±2.6	7.9±2.8	6.4±2.5	5.9±2.4	5.1±2.2	5.2±2.3	4.4±2.1	6.2±2.5
Central Michigan District	5.2±1.9	4.8±1.8	4.0±1.7	5.8±2.1	5.3±2.0	5.4±2.0	4.5±1.9	5.5±2.0	5.3±2.0	6.0±2.1
Chippewa County	*	*	*	*	*	*	*	*	*	*
City Of Detroit	5.2±1.9	5.5±2.1	5.8±2.3	6.5±2.8	10.3±3.9	10.1±4.0	11.3±4.2	9.5±3.8	9.2±3.7	8.4±3.4
Dickinson-Iron District	6.9±5.1	*	8.2±5.6	7.2±5.3	*	*	5.9±4.7	6.1±4.9	*	*
District #10	6.4±1.6	6.3±1.7	6.4±1.7	6.3±1.7	7.7±1.9	6.8±1.8	8.2±2.0	7.0±1.8	6.9±1.8	5.0±1.6
District #2	5.5±3.6	8.0±4.3	10.1±4.9	9.3±4.7	9.2±4.6	6.7±3.9	7.2±4.0	4.2±3.1	4.7±3.2	3.6±2.9
District #4	6.6±3.4	6.4±3.4	6.3±3.6	6.5±3.7	6.2±3.7	4.5±3.1	4.4±3.1	3.8±2.8	4.3±3.0	6.3±3.6
Genesee County	6.0±1.4	5.4±1.3	5.1±1.3	5.1±1.4	4.8±1.3	5.2±1.4	5.3±1.4	5.7±1.5	6.3±1.6	6.4±1.6
Grand Traverse County	7.4±3.2	7.3±3.2	6.0±3.0	3.5±2.3	3.1±2.2	*	3.7±2.3	5.6±2.8	7.8±3.3	6.8±3.1
Huron County	*	*	*	6.5±5.2	6.7±5.3	*	*	-	*	*
Ingham County	4.1±1.4	4.7±1.5	3.6±1.4	4.6±1.6	5.3±1.7	7.5±2.1	7.1±2.0	6.4±1.9	4.6±1.6	4.3±1.6
Ionia County	4.1±2.5	3.4±2.4	3.1±2.3	3.6±2.5	3.7±2.6	3.7±2.6	4.2±2.8	5.1±3.0	4.8±2.9	4.3±2.8
Jackson County	6.4±2.2	6.2±2.2	5.7±2.1	5.5±2.1	4.1±1.8	4.6±2.0	3.9±1.8	5.1±2.1	4.8±2.0	5.6±2.2
Kalamazoo County	4.8±1.6	5.5±1.7	5.7±1.7	4.8±1.6	3.9±1.4	3.9±1.4	4.7±1.6	4.7±1.6	3.2±1.3	4.7±1.6
Kent County	5.6±1.0	5.0±0.9	5.0±0.9	4.4±0.9	4.4±0.9	4.1±0.9	4.5±0.9	4.9±0.9	5.1±1.0	4.8±0.9
Lapeer County	5.5±2.8	7.3±3.3	7.8±3.5	6.3±3.2	3.0±2.2	2.6±2.1	4.7±2.8	5.6±3.0	5.6±3.0	4.8±2.9
Lenawee County	9.0±3.3	8.4±3.3	10.6±3.7	8.7±3.4	9.6±3.6	7.6±3.2	8.4±3.3	7.6±3.2	7.0±3.1	6.5±3.0
Livingston County	5.3±1.9	6.6±2.2	6.1±2.2	6.6±2.2	5.8±2.1	5.0±2.0	5.1±2.0	4.7±1.9	4.2±1.7	3.0±1.5
Luce-Mackinac-Alger-Schoolcraft	11.7±7.6	10.5±7.2	9.5±7.0	*	-	-	*	*	11.0±8.1	*
Macomb County	5.2±0.9	6.2±1.0	5.9±1.0	5.6±1.0	4.8±0.9	5.5±1.0	5.3±1.0	5.4±1.0	4.6±0.9	5.2±1.0
Marquette County	4.2±2.9	3.6±2.7	3.7±2.8	4.8±3.1	4.2±2.9	4.3±3.0	*	4.5±3.1	4.4±3.0	4.5±3.1
Midland County	5.6±2.9	4.0±2.5	3.7±2.4	4.0±2.5	4.9±2.8	5.8±3.0	6.3±3.2	4.9±2.8	3.7±2.4	2.5±2.0

**Table 19-B
White Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2006-2017**

Health Department	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Mid-Michigan District	5.0±1.8	4.7±1.8	5.6±1.9	5.8±2.0	5.4±1.9	4.3±1.7	4.9±1.8	5.5±2.0	5.4±1.9	4.4±1.8
Monroe County	6.4±2.3	4.8±2.0	4.7±1.9	3.5±1.7	3.6±1.8	5.1±2.1	5.3±2.2	6.0±2.3	4.1±1.9	6.1±2.4
Muskegon County	5.7±2.0	6.4±2.2	6.5±2.2	6.3±2.2	7.1±2.4	6.6±2.3	5.8±2.2	4.6±1.9	4.1±1.8	4.2±1.8
Northwest Michigan Community Health	6.4±2.7	6.5±2.8	6.4±2.9	6.8±3.0	6.9±3.0	6.8±3.0	5.8±2.8	5.7±2.8	3.5±2.2	2.6±1.9
Oakland County	4.9±0.8	4.8±0.8	5.3±0.8	5.1±0.8	5.7±0.9	5.6±0.9	5.4±0.8	4.5±0.8	4.0±0.7	3.8±0.7
Ottawa County	5.3±1.4	4.6±1.4	4.9±1.4	5.3±1.5	5.6±1.5	6.3±1.6	5.3±1.5	5.6±1.5	4.7±1.4	4.6±1.4
Delta & Menominee	3.8±3.0	4.3±3.4	*	*	4.5±3.3	3.8±3.0	4.9±3.4	*	*	*
Saginaw County	6.6±2.2	6.4±2.2	4.7±1.9	2.9±1.5	2.9±1.5	4.2±1.8	6.2±2.2	6.3±2.2	7.5±2.5	5.5±2.1
Sanilac County	7.0±4.3	6.3±4.1	*	*	*	4.6±3.7	6.9±4.5	6.1±4.2	7.7±4.8	7.8±4.8
Shiawassee County	5.5±3.0	5.9±3.2	3.3±2.4	3.4±2.5	3.9±2.7	3.3±2.5	4.3±2.8	4.3±2.8	5.8±3.3	6.9±3.6
St Clair County	5.0±1.9	6.2±2.1	5.4±2.1	6.3±2.3	6.2±2.3	6.3±2.3	5.9±2.3	4.3±1.9	5.0±2.1	4.7±2.1
Tuscola County	3.9±2.9	4.0±3.0	6.1±3.8	6.9±4.1	7.0±4.1	5.0±3.5	4.9±3.4	4.8±3.3	4.3±3.2	4.4±3.2
Van Buren-Cass County	5.0±2.2	5.9±2.4	6.0±2.4	4.5±2.1	5.8±2.5	6.7±2.7	7.1±2.7	6.3±2.5	6.9±2.6	6.2±2.5
Washtenaw County	3.5±1.3	4.3±1.4	4.4±1.5	4.5±1.5	3.9±1.4	4.2±1.5	5.4±1.7	5.0±1.6	4.3±1.5	3.2±1.3
Wayne County	5.6±0.8	5.6±0.8	5.4±0.8	5.4±0.8	5.5±0.9	6.0±0.9	6.0±0.9	6.1±0.9	5.8±0.9	5.4±0.8
Western Upper Peninsula District	7.3±3.8	4.8±3.1	4.7±3.1	3.8±2.8	4.4±3.1	4.1±3.0	5.3±3.5	4.8±3.3	5.9±3.7	6.6±3.9

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 19-C
Black Three-Year Moving Average Infant Death Rates by District
Michigan Residents, 2006-2017

Health Department	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	15.3±0.9	15.5±0.9	14.8±0.9	14.5±0.9	13.8±0.9	13.4±0.9	13.3±0.9	13.7±0.9	13.8±0.9	14.3±0.9
Barry-Eaton District	*	25.2±18.4	28.0±19.1	26.9±18.4	*	22.7±16.6	22.4±16.4	29.0±17.7	24.0±15.5	22.0±15.1
Berrien County	14.9±6.0	16.7±6.4	16.0±6.2	19.5±6.9	18.6±6.9	17.3±6.7	16.5±6.7	13.3±6.1	10.4±5.6	7.4±4.8
Calhoun County	24.2±10.5	20.5±9.6	22.1±10.4	25.4±11.3	20.2±10.1	13.8±8.1	9.9±6.8	14.6±8.2	11.3±7.3	12.6±7.8
City Of Detroit	15.8±1.4	16.5±1.5	16.0±1.5	15.0±1.5	14.7±1.5	14.8±1.5	15.0±1.5	14.8±1.5	14.6±1.5	14.8±1.5
District #10	*	*	*	*	*	*	*	*	-	*
Genesee County	17.1±3.6	17.9±3.7	14.6±3.4	14.2±3.4	12.4±3.3	14.2±3.5	14.6±3.6	15.6±3.7	16.5±3.9	15.2±3.7
Ingham County	17.8±6.3	16.5±6.0	16.7±5.9	13.1±5.3	11.4±5.0	10.1±4.7	10.1±4.7	12.9±5.2	14.2±5.5	14.5±5.6
Jackson County	16.8±10.3	15.6±10.1	13.2±9.1	11.6±8.6	13.0±9.0	18.5±10.8	16.2±10.0	15.0±9.7	20.2±11.3	26.5±12.8
Kalamazoo County	16.8±6.2	19.2±6.5	19.5±6.5	18.3±6.4	17.6±6.3	15.5±5.9	14.2±5.7	15.3±5.9	11.0±4.9	15.3±5.6
Kent County	14.4±3.8	13.4±3.7	13.9±3.8	16.6±4.2	14.6±3.9	12.9±3.6	9.9±3.2	10.7±3.3	13.0±3.7	12.8±3.6
Macomb County	13.4±4.0	15.1±4.0	13.8±3.8	13.1±3.6	11.4±3.2	11.1±3.1	9.9±2.9	10.7±2.9	10.6±2.9	12.8±3.1
Mid-Michigan District	-	-	-	*	*	*	-	*	*	*
Muskegon County	9.0±4.9	8.2±4.6	13.2±5.9	14.9±6.3	16.4±6.8	12.8±6.0	14.9±6.7	19.7±7.7	23.1±8.3	23.2±8.1
Oakland County	15.4±3.1	14.7±3.0	11.9±2.7	10.8±2.5	10.6±2.4	11.1±2.5	13.7±2.8	14.7±2.9	14.3±2.9	13.5±2.8
Ottawa County	*	*	*	*	*	*	*	*	*	*
Saginaw County	15.3±5.2	19.2±5.8	17.4±5.6	19.2±6.1	13.4±5.1	14.3±5.2	12.2±5.0	15.9±5.7	17.3±5.9	16.2±5.6
St Clair County	*	*	*	*	*	*	*	*	34.3±25.0	33.7±24.5
Van Buren-Cass County	*	*	*	*	*	*	*	*	*	*
Washtenaw County	13.2±4.9	11.6±4.6	10.5±4.4	8.7±4.0	8.9±4.1	8.3±4.1	10.9±4.7	12.1±5.2	16.4±5.9	14.4±5.5
Wayne County	12.7±2.7	10.4±2.4	11.7±2.5	12.6±2.5	13.6±2.6	11.3±2.4	9.6±2.2	8.1±2.0	9.0±2.1	12.1±2.4

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

-: A dash indicates that there were either no births or no infant deaths from this category.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 20-A
Three-Year Moving Average Infant Death Rates by City
Michigan Residents, 2006-2017

City	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	7.6±0.3	7.6±0.3	7.3±0.3	7.1±0.3	6.9±0.3	6.8±0.3	6.9±0.3	6.9±0.3	6.7±0.3	6.7±0.3
Ann Arbor	3.9±2.1	4.3±2.2	3.7±2.1	3.1±1.9	2.7±1.8	3.6±2.0	5.1±2.4	4.3±2.2	3.7±2.1	3.2±2.0
Battle Creek	12.4±4.3	11.0±4.1	10.6±4.1	11.3±4.3	11.0±4.3	9.9±3.8	7.9±3.4	8.4±3.5	7.0±3.4	9.6±4.1
Dearborn	7.7±2.3	6.9±2.3	7.0±2.3	6.3±2.2	7.4±2.4	7.5±2.4	6.4±2.2	6.0±2.1	4.9±1.9	5.5±2.0
Dearborn Heights	5.8±3.0	5.8±3.0	3.8±2.5	5.2±2.9	5.7±3.1	6.9±3.4	5.5±3.0	5.2±2.8	6.7±3.2	5.5±2.9
Detroit	14.3±1.2	14.8±1.3	14.4±1.3	13.6±1.3	13.6±1.3	13.6±1.3	13.9±1.3	13.6±1.3	13.6±1.3	14.0±1.4
Farmington Hills	7.4±3.3	7.0±3.2	6.4±3.1	6.4±3.1	6.9±3.3	6.7±3.2	6.7±3.2	6.1±3.0	4.5±2.5	6.2±2.9
Flint	11.6±2.7	12.6±2.9	12.6±3.0	12.1±3.0	11.6±3.0	11.0±2.8	10.9±2.8	10.7±2.9	13.1±3.4	13.9±3.5
Grand Rapids	10.0±1.9	8.3±1.8	8.6±1.8	8.5±1.8	9.4±1.9	8.0±1.7	6.6±1.6	5.9±1.5	6.8±1.7	7.3±1.7
Kalamazoo	9.2±3.3	10.5±3.5	11.0±3.6	10.3±3.6	9.7±3.5	6.0±2.5	6.0±2.5	7.6±2.8	7.3±3.0	8.4±3.3
Kentwood	4.2±2.6	3.9±2.6	3.4±2.4	3.4±2.4	4.5±2.7	5.2±2.9	5.1±2.9	6.0±3.1	8.1±3.5	8.9±3.7
Lansing	8.5±2.3	8.9±2.4	8.7±2.4	7.1±2.2	6.5±2.1	7.4±2.2	8.5±2.4	8.8±2.4	9.2±2.5	8.7±2.5
Livonia	5.4±2.8	5.2±2.8	4.8±2.7	2.4±1.9	2.8±2.1	5.5±2.9	5.4±2.8	4.8±2.6	4.0±2.4	5.2±2.7
Midland	5.9±4.1	5.1±3.8	*	*	*	6.3±4.1	7.7±4.5	7.7±4.5	4.2±3.4	*
Muskegon	7.5±3.9	7.5±3.9	7.2±3.9	7.8±4.1	10.5±4.8	12.6±5.0	12.6±5.0	13.4±5.2	14.9±5.9	13.9±5.8
Novi	5.3±3.3	5.3±3.3	6.6±3.7	3.3±2.7	3.8±2.8	4.3±3.0	6.1±3.5	5.0±3.1	4.5±2.9	3.7±2.7
Pontiac	12.5±3.7	14.4±4.1	13.8±4.0	11.9±3.8	8.5±3.2	9.4±3.3	11.4±3.6	14.0±4.0	12.6±3.8	11.8±3.7
Portage	4.4±3.1	9.2±4.5	8.6±4.3	7.5±4.1	4.1±3.0	5.8±3.6	5.8±3.6	5.2±3.4	*	7.7±4.2
Rochester Hills	6.2±3.1	5.0±2.8	4.8±2.8	3.5±2.5	4.0±2.6	5.2±3.0	4.4±2.7	4.0±2.6	2.7±2.1	2.7±2.1
Roseville	9.7±4.4	10.0±4.6	5.5±3.4	5.7±3.5	7.4±4.0	7.8±4.1	7.8±4.0	7.7±4.0	10.1±4.7	10.2±4.7
Royal Oak	4.0±2.6	4.4±2.7	5.6±3.0	6.4±3.2	8.2±3.7	7.1±3.5	7.6±3.6	4.0±2.6	2.6±2.1	*
Saginaw	11.5±3.8	14.6±4.3	13.2±4.2	13.7±4.4	9.6±3.7	12.1±4.2	12.7±4.3	15.1±4.8	15.8±5.0	13.4±4.6
Southfield	12.4±4.5	13.1±4.6	11.2±4.4	10.0±4.1	10.9±4.3	8.5±3.7	11.7±4.4	9.0±3.8	9.8±4.0	7.6±3.5
St Clair Shores	3.7±2.7	5.4±3.4	4.0±3.0	4.7±3.3	*	4.7±3.2	4.6±3.2	5.6±3.5	6.3±3.7	5.8±3.6
Sterling Heights	7.0±2.4	7.5±2.5	7.0±2.5	6.2±2.4	6.6±2.4	5.7±2.3	6.7±2.4	6.4±2.4	6.9±2.5	6.7±2.4
Taylor	8.5±3.5	6.3±3.1	6.8±3.2	9.9±3.9	11.8±4.2	10.8±4.0	7.4±3.3	6.2±3.0	6.5±3.1	7.8±3.4
Troy	5.8±3.0	5.5±3.0	6.2±3.2	4.4±2.7	3.8±2.5	3.3±2.3	3.8±2.4	4.1±2.5	4.1±2.5	4.5±2.6
Warren	6.4±2.2	8.1±2.5	7.9±2.5	9.1±2.7	7.6±2.4	8.7±2.6	6.6±2.2	5.4±2.0	5.3±2.0	6.5±2.2
Westland	7.5±2.9	7.4±2.9	8.2±3.1	9.6±3.4	10.4±3.5	9.3±3.3	8.3±3.1	7.4±2.9	7.3±2.9	6.5±2.8
Wyoming	12.1±3.4	11.0±3.3	8.4±3.0	6.7±2.6	4.6±2.2	4.1±2.1	4.4±2.1	6.0±2.5	6.6±2.6	5.9±2.5

Notes: This table shows cities with (1) populations greater than 40,000 according to 2007 estimates and (2) three-year average births of greater than 200. City birth and mortality statistics have changed from pre-2011 reports due to geocoding. See Data Quality Considerations for further details. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 20-B
White Three-Year Moving Average Infant Death Rates by City
Michigan Residents, 2006-2017

City	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	5.6±0.3	5.6±0.3	5.4±0.3	5.2±0.3	5.3±0.3	5.3±0.3	5.5±0.3	5.4±0.3	5.1±0.3	4.9±0.3
Ann Arbor	*	3.7±2.5	3.2±2.4	3.7±2.6	*	3.2±2.3	4.5±2.8	4.0±2.6	2.8±2.2	*
Battle Creek	9.5±4.4	7.2±3.9	4.8±3.3	5.5±3.6	8.1±4.4	8.1±4.1	7.0±3.8	5.5±3.4	4.6±3.4	9.0±4.9
Dearborn	7.4±2.4	6.4±2.3	6.2±2.3	6.0±2.3	6.5±2.4	7.0±2.5	5.9±2.4	7.0±2.6	5.6±2.3	5.4±2.1
Dearborn Heights	5.2±3.1	4.3±2.8	3.0±2.4	3.1±2.5	3.7±2.7	5.3±3.2	6.0±3.5	6.9±3.8	8.7±4.1	6.3±3.4
Detroit	5.2±1.9	5.5±2.1	5.8±2.3	6.5±2.8	9.9±3.8	10.1±4.0	11.3±4.2	9.9±3.9	9.2±3.7	8.4±3.4
Farmington Hills	7.2±4.3	6.1±4.0	5.6±3.9	6.5±4.2	8.0±4.7	6.5±4.2	5.0±3.7	4.9±3.6	4.1±3.3	4.1±3.3
Flint	5.7±3.1	6.7±3.5	9.6±4.3	7.1±3.8	7.6±4.1	5.5±3.4	5.9±3.5	4.5±3.1	7.1±4.2	7.7±4.5
Grand Rapids	5.5±1.7	4.7±1.6	5.1±1.7	4.4±1.6	5.7±1.8	5.3±1.7	4.6±1.6	4.4±1.6	4.5±1.7	5.0±1.8
Kalamazoo	3.6±2.7	*	3.6±2.7	4.8±3.1	6.1±3.6	3.5±2.4	2.6±2.0	3.0±2.2	*	3.4±2.7
Kentwood	*	*	*	*	*	4.5±3.3	5.2±3.6	5.4±3.7	7.0±4.1	7.9±4.4
Lansing	4.5±2.1	5.2±2.3	4.6±2.2	5.2±2.4	5.3±2.5	7.4±2.9	7.9±3.1	7.6±3.1	6.3±2.9	5.2±2.7
Livonia	5.6±3.0	5.4±3.0	5.0±2.9	*	*	4.9±2.9	5.3±3.0	5.1±2.9	3.8±2.5	4.2±2.6
Midland	6.4±4.4	5.5±4.1	*	*	*	4.7±3.7	6.1±4.2	6.1±4.2	4.6±3.6	*
Muskegon	7.2±4.9	6.7±4.9	*	*	7.1±5.2	8.6±5.3	6.0±4.4	5.3±4.2	*	*
Novi	*	*	5.1±4.1	*	*	*	5.1±4.1	*	*	*
Pontiac	5.7±3.5	8.5±4.6	7.7±4.6	6.3±4.4	*	4.9±3.9	4.8±3.8	6.2±4.3	6.4±4.4	6.8±4.7
Portage	3.8±3.0	6.8±4.2	7.4±4.4	6.1±4.0	4.1±3.3	5.4±3.7	5.4±3.7	4.7±3.5	*	6.2±4.0
Rochester Hills	7.0±3.8	5.5±3.4	5.6±3.5	3.4±2.7	3.4±2.7	3.9±2.9	*	3.5±2.8	*	*
Roseville	9.8±4.9	10.3±5.2	6.3±4.1	5.3±3.9	5.6±4.1	4.7±3.8	6.3±4.3	5.5±4.1	9.0±5.3	8.5±5.2
Royal Oak	3.0±2.4	3.4±2.5	5.8±3.3	6.8±3.5	8.3±3.9	6.5±3.5	6.9±3.6	3.9±2.7	2.9±2.3	*
Saginaw	6.2±4.0	6.3±4.1	4.3±3.5	*	*	5.7±4.2	9.7±5.4	11.4±5.9	13.0±6.5	8.2±5.4
Southfield	*	*	11.6±9.2	11.9±9.5	12.0±9.5	*	*	-	*	*
St Clair Shores	4.1±3.0	6.0±3.7	4.5±3.4	5.3±3.6	*	*	*	4.9±3.4	6.3±3.9	5.2±3.6
Sterling Heights	6.1±2.5	6.9±2.7	7.7±2.9	6.9±2.8	7.6±3.0	6.0±2.6	6.5±2.7	5.4±2.4	5.7±2.5	5.7±2.5
Taylor	7.0±3.8	5.6±3.5	5.3±3.5	6.7±4.0	8.2±4.4	8.0±4.3	6.8±4.0	6.1±3.8	6.2±3.8	5.2±3.6
Troy	5.1±3.5	5.3±3.6	6.1±4.0	5.4±3.7	5.9±3.9	4.6±3.4	4.7±3.4	3.9±3.1	*	4.1±3.3
Warren	5.9±2.5	6.8±2.7	6.3±2.7	6.9±2.9	5.9±2.7	8.0±3.2	6.7±2.9	5.7±2.7	4.7±2.5	5.5±2.7
Westland	4.9±2.8	5.9±3.1	5.7±3.1	7.8±3.7	8.6±3.9	8.5±3.9	9.0±4.0	8.0±3.8	7.2±3.6	4.5±2.9
Wyoming	9.3±3.2	8.6±3.2	6.7±2.8	5.3±2.5	3.8±2.1	3.2±2.0	4.2±2.3	5.7±2.7	5.5±2.7	4.3±2.4

Notes: This table shows cities with (1) populations greater than 40,000 according to 2007 estimates and (2) three-year average births of greater than 200. City birth and mortality statistics have changed from pre-2011 reports due to geocoding. See Data Quality Considerations for further details. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 20-C
Black Three-Year Moving Average Infant Death Rates by City
Michigan Residents, 2006-2017

City	2006-08	2007-09	2008-10	2009-11	2010-12	2011-13	2012-14	2013-15	2014-16	2015-17
Michigan	15.3±0.9	15.5±0.9	14.8±0.9	14.5±0.9	13.8±0.9	13.4±0.9	13.3±0.9	13.7±0.9	13.8±0.9	14.3±0.9
Battle Creek	20.9±11.7	21.0±11.7	23.0±12.9	21.5±12.6	11.9±9.5	12.1±8.9	11.6±8.5	16.0±9.9	10.2±8.1	*
Detroit	15.8±1.4	16.5±1.5	15.9±1.5	15.0±1.5	14.7±1.5	14.8±1.5	15.0±1.5	14.8±1.5	14.5±1.5	14.7±1.5
Flint	15.5±4.1	16.6±4.2	14.7±4.1	15.3±4.3	14.1±4.2	14.7±4.2	14.4±4.2	14.8±4.3	16.4±4.8	16.8±4.9
Grand Rapids	17.6±5.1	16.8±5.0	16.0±5.0	18.8±5.4	16.7±5.1	14.9±4.8	11.5±4.2	12.0±4.4	13.9±4.9	13.1±4.6
Kalamazoo	18.5±8.0	22.4±8.7	21.6±8.5	17.7±7.9	12.7±6.9	9.7±5.7	11.5±6.2	17.0±7.6	14.6±7.3	14.6±7.1
Lansing	19.0±6.7	17.9±6.4	18.3±6.5	13.5±5.7	11.0±5.2	8.9±4.7	10.7±5.1	13.6±5.6	16.2±6.2	16.0±6.1
Muskegon	*	*	12.0±8.3	13.1±8.5	17.4±10.2	18.1±10.1	20.1±10.8	21.6±11.2	29.1±13.3	26.5±12.4
Pontiac	19.6±6.7	19.9±6.8	19.1±6.7	16.5±6.2	12.5±5.3	12.6±5.2	16.4±5.9	19.4±6.5	17.9±6.2	16.1±5.9
Saginaw	16.0±6.2	20.1±7.0	18.4±6.9	20.5±7.5	14.1±6.3	15.8±6.5	14.2±6.3	17.5±7.2	18.1±7.5	17.5±7.2
Southfield	14.8±5.8	15.0±5.8	11.6±5.2	10.0±4.7	11.2±5.0	10.3±4.7	14.9±5.8	12.4±5.3	13.2±5.5	10.2±4.8
Taylor	14.9±9.7	9.3±7.4	11.9±8.2	18.8±9.8	21.6±10.2	18.7±9.4	10.2±7.0	7.6±6.0	7.2±5.7	9.6±6.6
Warren	11.8±7.2	14.8±7.7	14.6±7.3	14.1±6.9	11.9±6.0	9.4±5.1	7.8±4.6	5.6±3.9	6.8±4.2	9.4±4.9
Westland	12.3±8.5	11.0±7.6	14.0±8.2	12.1±7.5	14.7±8.3	11.7±7.2	9.1±6.3	6.6±5.3	8.6±5.9	9.4±6.1

Notes: This table shows cities with (1) populations greater than 40,000 according to 2007 estimates and (2) three-year average births of greater than 200. City birth and mortality statistics have changed from pre-2011 reports due to geocoding. See Data Quality Considerations for further details. Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

*: A rate is not calculated when there are fewer than 6 events, because the width of the confidence interval would negate any usefulness for comparative purposes.

Source: 2006-2017 Geocoded Michigan Death Certificate Registries 2006-2017 Geocoded Michigan Birth Certificate Registries. Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 21-A
Number of Neonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2007-2017

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Michigan	697	608	600	549	501	540	544	513	498	491	506
Alcona	-	1	1	-	-	1	-	1	-	-	1
Alger	-	3	-	-	-	-	-	-	-	-	-
Allegan	13	4	4	6	5	3	3	4	9	3	3
Alpena	-	1	-	-	1	-	1	1	1	-	1
Antrim	-	2	-	4	2	-	3	-	-	-	-
Arenac	-	-	2	-	1	1	4	2	-	-	-
Baraga	-	1	-	-	-	-	1	1	1	-	1
Barry	3	2	3	4	2	2	3	1	-	2	4
Bay	5	3	7	9	6	4	-	5	4	1	5
Benzie	2	-	-	-	-	-	2	-	1	2	1
Berrien	13	10	10	8	8	11	8	9	6	2	2
Branch	2	1	2	2	3	3	2	3	2	4	4
Calhoun	18	13	13	10	9	7	8	5	8	4	12
Cass	2	2	5	1	2	2	-	5	-	3	3
Charlevoix	2	1	2	-	-	-	-	-	2	-	1
Cheboygan	1	-	-	-	-	-	-	-	-	-	2
Chippewa	-	1	-	-	3	2	1	1	-	1	1
Clare	3	-	1	-	1	2	1	1	-	1	1
Clinton	6	1	1	3	2	2	1	3	5	1	2
Crawford	4	1	1	-	-	-	1	-	-	3	-
Delta	2	-	2	2	-	4	1	3	-	-	1
Dickinson	-	2	2	1	-	-	4	1	-	-	-
Eaton	4	3	3	7	5	2	6	3	7	7	2
Emmet	1	2	-	3	-	1	2	1	-	1	-
Genesee	40	27	34	19	22	19	25	23	29	31	15

Table 21-A
Number of Neonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2007-2017

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Gladwin	1	1	1	1	1	-	1	1	2	1	-
Gogebic	1	-	-	2	-	-	-	1	-	3	-
Grand Traverse	5	7	3	3	-	3	-	1	7	4	3
Gratiot	4	2	3	3	3	-	3	2	-	-	2
Hillsdale	4	3	1	1	2	1	3	-	3	4	5
Houghton	2	3	1	3	1	2	2	-	-	2	3
Huron	-	-	3	1	1	-	-	-	-	1	1
Ingham	17	22	14	12	16	16	20	15	11	5	8
Ionia	2	-	1	2	1	-	2	2	2	2	1
Iosco	-	-	1	1	1	-	-	2	1	-	2
Iron	1	-	-	1	-	-	-	-	-	-	1
Isabella	5	1	1	-	6	2	-	2	4	-	6
Jackson	12	11	7	9	8	3	10	5	7	12	7
Kalamazoo	22	13	20	17	8	17	11	11	11	5	25
Kalkaska	-	-	1	1	-	3	1	3	-	2	-
Kent	47	48	42	32	40	38	28	37	38	45	31
Keweenaw	-	-	-	-	-	-	-	-	-	-	-
Lake	-	-	1	-	-	1	-	-	1	-	1
Lapeer	4	3	6	4	-	1	4	3	1	4	3
Leelanau	-	-	-	3	-	-	-	1	-	1	-
Lenawee	5	9	8	11	3	6	4	5	4	5	3
Livingston	11	6	9	6	9	7	3	9	5	3	7
Luce	-	1	-	-	-	-	-	1	-	-	-
Mackinac	-	1	1	-	-	-	-	-	2	3	-
Macomb	48	40	52	39	43	40	52	34	43	38	52

Table 21-A
Number of Neonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2007-2017

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Manistee	-	1	1	1	-	-	-	1	-	-	1
Marquette	2	2	2	1	3	-	2	-	1	1	-
Mason	-	1	1	2	1	-	-	1	2	-	-
Mecosta	1	4	-	1	4	4	4	2	3	-	1
Menominee	1	1	1	-	1	-	-	1	1	-	-
Midland	3	1	3	2	2	3	4	3	-	1	2
Missaukee	-	-	1	-	-	1	-	2	-	-	-
Monroe	6	5	3	4	2	6	10	2	8	3	10
Montcalm	2	1	2	5	3	3	4	1	2	3	-
Montmorency	-	-	-	1	-	-	-	-	-	-	-
Muskegon	13	9	9	16	11	11	14	9	7	15	11
Newaygo	6	3	1	2	6	2	2	3	-	3	3
Oakland	71	62	60	63	45	61	69	66	53	55	55
Oceana	1	2	2	3	-	2	1	3	1	3	-
Ogemaw	1	-	-	1	1	1	-	-	-	-	-
Ontonagon	1	-	-	-	-	-	1	1	-	-	-
Osceola	1	-	1	1	-	-	1	-	2	2	1
Oscoda	-	1	1	-	-	1	-	1	-	-	-
Otsego	3	1	-	2	3	1	-	1	1	1	-
Ottawa	16	19	10	17	17	14	20	10	12	9	9
Presque Isle	2	-	-	1	1	3	-	1	2	1	2
Roscommon	3	1	1	-	1	-	-	1	1	-	-
Saginaw	16	10	15	10	8	8	13	11	12	16	4
St. Clair	10	4	6	4	6	10	2	3	6	7	4
St. Joseph	6	8	9	2	3	1	5	3	1	5	4

Table 21-A
Number of Neonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2007-2017

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Sanilac	4	1	2	-	-	1	1	3	3	4	3
Schoolcraft	1	-	-	-	-	-	-	-	-	-	-
Shiawassee	8	-	2	2	1	2	2	1	1	2	4
Tuscola	1	-	2	4	1	4	1	-	3	-	1
Van Buren	5	6	4	2	3	4	4	3	3	5	1
Washtenaw	17	14	18	11	10	14	12	23	10	15	10
Wayne	180	197	172	158	151	172	147	147	144	129	155
Detroit City	120	123	114	98	80	104	87	80	81	76	86
Wayne County Excluding Detroit City	60	74	58	60	71	68	60	67	63	53	69
Wexford	4	2	2	2	2	5	3	2	1	1	2
Unknown	-	-	-	-	-	-	1	-	1	4	-

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Neonatal Death: Deaths occurring to individuals less than 28 days of age.
 -: A dash indicates quantity zero.

Source: 2007-2017 Michigan Resident Death Files,
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

**Table 21-B
Number of Postneonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2007-2017**

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Michigan	300	286	281	268	248	243	255	260	267	256	256
Alcona	-	-	-	-	1	-	-	-	-	-	-
Alger	-	-	1	-	-	-	-	-	-	-	-
Allegan	5	2	2	-	2	1	6	5	1	2	1
Alpena	1	-	3	1	1	-	-	1	-	-	1
Antrim	2	-	1	-	-	-	-	-	-	-	-
Arenac	-	-	-	1	1	-	-	-	-	1	-
Baraga	-	-	-	-	-	-	-	-	-	1	-
Barry	2	3	2	4	-	1	1	1	-	-	1
Bay	3	5	1	3	3	1	2	5	2	-	2
Benzie	-	-	-	1	-	-	-	-	-	-	-
Berrien	8	6	7	6	10	3	6	6	3	8	6
Branch	-	1	-	2	1	1	1	1	1	1	1
Calhoun	10	1	7	8	6	3	3	4	6	2	3
Cass	1	2	1	1	1	2	3	-	3	-	-
Charlevoix	-	-	-	-	1	-	-	1	1	-	1
Cheboygan	1	1	2	-	-	-	-	-	1	1	-
Chippewa	1	-	-	2	1	-	1	1	-	-	-
Clare	-	1	1	1	-	-	-	-	1	1	2
Clinton	-	2	1	2	-	-	1	2	3	2	-
Crawford	1	-	-	-	1	-	-	-	2	-	1
Delta	2	-	1	-	1	-	-	-	1	2	-
Dickinson	-	-	-	1	-	-	-	-	-	2	-
Eaton	2	4	3	1	1	3	4	3	2	2	1
Emmet	-	-	1	-	-	1	-	-	-	-	1
Genesee	17	19	17	11	18	19	15	18	14	17	22

**Table 21-B
Number of Postneonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2007-2017**

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Gladwin	-	-	1	1	-	-	1	2	2	-	-
Gogebic	-	-	-	-	-	-	-	-	-	2	-
Grand Traverse	2	1	2	-	1	1	-	5	2	2	-
Gratiot	-	1	-	2	-	-	-	2	1	1	-
Hillsdale	1	1	-	1	-	1	-	-	2	3	1
Houghton	1	1	-	-	-	-	1	1	-	-	2
Huron	-	-	-	-	1	3	-	-	-	-	2
Ingham	9	3	6	2	5	6	8	6	10	16	8
Ionia	4	1	-	3	2	1	3	1	1	2	1
Iosco	1	1	2	2	1	1	1	-	-	1	2
Iron	-	-	1	1	1	-	-	1	-	-	-
Isabella	2	-	3	4	-	2	2	-	1	1	2
Jackson	3	3	5	2	3	4	5	3	3	5	7
Kalamazoo	7	9	9	13	3	2	15	5	9	3	10
Kalkaska	-	-	-	1	-	1	1	-	-	-	-
Kent	18	21	14	22	19	12	11	15	18	11	15
Keweenaw	-	-	-	-	-	-	-	-	-	-	-
Lake	-	-	1	1	-	-	-	-	-	-	1
Lapeer	1	1	5	3	-	1	-	2	3	-	-
Leelanau	1	-	-	-	1	-	1	2	1	-	-
Lenawee	4	2	1	4	3	2	5	4	2	3	2
Livingston	3	4	3	5	3	1	4	4	2	1	1
Luce	-	1	-	-	-	-	-	1	1	1	-
Mackinac	-	1	-	-	-	-	-	-	-	-	-
Macomb	21	22	26	9	10	12	15	14	17	15	16

**Table 21-B
Number of Postneonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2007-2017**

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Manistee	-	-	1	1	1	-	-	1	-	1	2
Marquette	-	-	2	1	1	2	-	1	4	1	1
Mason	1	1	-	1	-	-	-	1	-	-	-
Mecosta	1	5	-	2	2	-	1	1	-	1	-
Menominee	-	1	-	-	-	1	-	-	-	1	-
Midland	-	2	2	-	2	3	3	2	3	-	-
Missaukee	-	-	-	-	-	-	-	-	-	1	-
Monroe	4	6	6	5	1	3	5	1	1	4	1
Montcalm	-	2	2	1	1	2	1	1	1	1	3
Montmorency	-	-	-	-	-	2	-	-	-	-	-
Muskegon	6	7	6	7	4	7	5	6	11	5	4
Newaygo	3	-	2	1	1	-	1	4	3	-	1
Oakland	21	26	17	21	25	32	21	17	18	17	16
Oceana	1	1	-	1	-	3	-	3	1	-	-
Ogemaw	1	1	1	1	-	1	-	1	-	1	1
Ontonagon	-	-	-	-	-	-	-	-	-	-	-
Osceola	1	-	-	1	2	-	-	-	-	-	1
Oscoda	-	1	-	-	-	1	-	-	-	-	-
Otsego	2	-	1	-	1	2	3	2	-	-	1
Ottawa	1	4	5	3	4	4	6	2	9	7	2
Presque Isle	-	-	1	1	-	-	-	-	-	-	-
Roscommon	1	-	-	-	-	-	-	-	-	1	-
Saginaw	10	15	11	2	9	5	6	10	7	11	6
St. Clair	2	4	9	2	5	3	4	6	3	4	3
St. Joseph	4	2	1	5	1	6	2	2	-	1	-

**Table 21-B
Number of Postneonatal Deaths by County of Residence,
State of Michigan, Michigan Counties and Detroit City
Michigan Residents, 2007-2017**

County	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Sanilac	2	-	1	1	-	3	1	-	1	-	-
Schoolcraft	-	-	-	-	-	-	-	-	-	-	-
Shiawassee	2	1	-	2	-	1	1	2	3	4	1
Tuscola	1	1	2	1	1	-	1	2	1	1	1
Van Buren	1	4	1	2	1	4	2	-	3	2	2
Washtenaw	6	3	4	9	4	6	6	5	4	7	5
Wayne	94	78	76	75	79	66	70	71	78	76	90
Detroit City	59	52	52	48	50	47	47	50	58	49	51
Wayne County Excluding Detroit City	35	26	24	27	29	19	23	21	20	27	39
Wexford	1	1	-	1	1	1	-	2	-	-	2
Unknown	-	-	-	-	-	-	-	1	-	-	-

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Postneonatal Death: Deaths occurring to individuals 28–364 days of age.
-: A dash indicates quantity zero.

Source: 2007-2017 Michigan Resident Death Files,
Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

Table 22-A
Number of Neonatal Deaths by Local Health Department and
State of Michigan,
Michigan Residents, 2007-2017

Health Department	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Michigan	697	608	600	549	501	540	544	513	498	491	506
Allegan County	13	4	4	6	5	3	3	4	9	3	3
Barry-Eaton District	7	5	6	11	7	4	9	4	7	9	6
Berrien County	13	10	10	8	8	11	8	9	6	2	2
Branch-Hillsdale-St. Joseph	12	12	12	5	8	5	10	6	6	13	13
Calhoun County	18	13	13	10	9	7	8	5	8	4	12
Central Michigan District	13	3	7	2	10	5	7	7	9	4	8
Chippewa County	-	1	-	-	3	2	1	1	-	1	1
City Of Detroit	120	123	114	98	80	104	87	80	81	76	86
Delta & Menominee	3	1	3	2	1	4	1	4	1	-	1
Dickinson-Iron District	1	2	2	2	-	-	4	1	-	-	1
District #10	16	14	11	12	13	18	12	17	8	12	8
District #2	1	2	3	2	2	3	-	4	1	-	3
District #4	3	1	-	2	2	3	1	2	3	1	5
Genesee County	40	27	34	19	22	19	25	23	29	31	15
Grand Traverse County	5	7	3	3	-	3	-	1	7	4	3
Ingham County	17	22	14	12	16	16	20	15	11	5	8
Ionia County	2	-	1	2	1	-	2	2	2	2	1
Jackson County	12	11	7	9	8	3	10	5	7	12	7
Kalamazoo County	22	13	20	17	8	17	11	11	11	5	25
Kent County	47	48	42	32	40	38	28	37	38	45	31
Lapeer County	4	3	6	4	-	1	4	3	1	4	3
Lenawee County	5	9	8	11	3	6	4	5	4	5	3
Livingston County	11	6	9	6	9	7	3	9	5	3	7
Luce-Mackinac-Alger-Schoolcraft	1	5	1	-	-	-	-	1	2	3	-
Macomb County	48	40	52	39	43	40	52	34	43	38	52

Table 22-A
Number of Neonatal Deaths by Local Health Department and
State of Michigan,
Michigan Residents, 2007-2017

Health Department	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Marquette County	2	2	2	1	3	-	2	-	1	1	-
Mid-Michigan District	12	4	6	11	8	5	8	6	7	4	4
Midland County	3	1	3	2	2	3	4	3	-	1	2
Monroe County	6	5	3	4	2	6	10	2	8	3	10
Muskegon County	13	9	9	16	11	11	14	9	7	15	11
Oakland County	71	62	60	63	45	61	69	66	53	55	55
Ottawa County	16	19	10	17	17	14	20	10	12	9	9
Bay County	5	3	7	9	6	4	-	5	4	1	5
Saginaw County	16	10	15	10	8	8	13	11	12	16	4
Shiawassee County	8	-	2	2	1	2	2	1	1	2	4
St Clair County	10	4	6	4	6	10	2	3	6	7	4
Van Buren-Cass County	7	8	9	3	5	6	4	8	3	8	4
Washtenaw County	17	14	18	11	10	14	12	23	10	15	10
Wayne County	60	74	58	60	71	68	60	67	63	53	69
Benzie-Leelanau District	2	-	-	3	-	-	2	1	1	3	1
Northwest Michigan Community Health	6	6	2	9	5	2	5	2	3	2	1
Western Upper Peninsula District	4	4	1	5	1	2	4	3	1	5	4
Tuscola County	1	-	2	4	1	4	1	-	3	-	1
Huron County	-	-	3	1	1	-	-	-	-	1	1
Sanilac County	4	1	2	-	-	1	1	3	3	4	3
Unknown	-	-	-	-	-	-	1	-	1	4	-

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Neonatal Death: Deaths occurring to individuals less than 28 days of age.
 -: A dash indicates quantity zero.

Source: 2007-2017 Michigan Resident Death Files,
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services

**Table 22-B
Number of Postneonatal Deaths by Local Health Department and
State of Michigan,
Michigan Residents, 2007-2017**

Health Department	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Michigan	300	286	281	268	248	243	255	260	267	256	256
Allegan County	5	2	2	–	2	1	6	5	1	2	1
Barry–Eaton District	4	7	5	5	1	4	5	4	2	2	2
Berrien County	8	6	7	6	10	3	6	6	3	8	6
Branch–Hillsdale–St. Joseph	5	4	1	8	2	8	3	3	3	5	2
Calhoun County	10	1	7	8	6	3	3	4	6	2	3
Central Michigan District	4	1	5	8	3	2	3	2	4	4	5
Chippewa County	1	–	–	2	1	–	1	1	–	–	–
City Of Detroit	59	52	52	48	50	47	47	50	58	49	51
Delta & Menominee	2	1	1	–	1	1	–	–	1	3	–
Dickinson–Iron District	–	–	1	2	1	–	–	1	–	2	–
District #10	8	8	4	9	6	5	3	12	6	3	7
District #2	2	3	3	3	2	3	1	1	–	2	3
District #4	2	1	6	2	1	2	–	1	1	1	1
Genesee County	17	19	17	11	18	19	15	18	14	17	22
Grand Traverse County	2	1	2	–	1	1	–	5	2	2	–
Ingham County	9	3	6	2	5	6	8	6	10	16	8
Ionia County	4	1	–	3	2	1	3	1	1	2	1
Jackson County	3	3	5	2	3	4	5	3	3	5	7
Kalamazoo County	7	9	9	13	3	2	15	5	9	3	10
Kent County	18	21	14	22	19	12	11	15	18	11	15
Lapeer County	1	1	5	3	–	1	–	2	3	–	–
Lenawee County	4	2	1	4	3	2	5	4	2	3	2
Livingston County	3	4	3	5	3	1	4	4	2	1	1
Luce–Mackinac–Alger–Schoolcraft	–	2	1	–	–	–	–	1	1	1	–
Macomb County	21	22	26	9	10	12	15	14	17	15	16

Table 22-B
Number of Postneonatal Deaths by Local Health Department and
State of Michigan,
Michigan Residents, 2007-2017

Health Department	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Marquette County	-	-	2	1	1	2	-	1	4	1	1
Mid-Michigan District	-	5	3	5	1	2	2	5	5	4	3
Midland County	-	2	2	-	2	3	3	2	3	-	-
Monroe County	4	6	6	5	1	3	5	1	1	4	1
Muskegon County	6	7	6	7	4	7	5	6	11	5	4
Oakland County	21	26	17	21	25	32	21	17	18	17	16
Ottawa County	1	4	5	3	4	4	6	2	9	7	2
Bay County	3	5	1	3	3	1	2	5	2	-	2
Saginaw County	10	15	11	2	9	5	6	10	7	11	6
Shiawassee County	2	1	-	2	-	1	1	2	3	4	1
St Clair County	2	4	9	2	5	3	4	6	3	4	3
Van Buren-Cass County	2	6	2	3	2	6	5	-	6	2	2
Washtenaw County	6	3	4	9	4	6	6	5	4	7	5
Wayne County	35	26	24	27	29	19	23	21	20	27	39
Benzie-Leelanau District	1	-	-	1	1	-	1	2	1	-	-
Northwest Michigan Community Health	4	-	3	-	2	3	3	3	1	-	3
Western Upper Peninsula District	1	1	-	-	-	-	1	1	-	3	2
Tuscola County	1	1	2	1	1	-	1	2	1	1	1
Huron County	-	-	-	-	1	3	-	-	-	-	2
Sanilac County	2	-	1	1	-	3	1	-	1	-	-
Unknown	-	-	-	-	-	-	-	1	-	-	-

Notes: Rates are per 1,000 live births. Adding and subtracting the number shown after the ± symbol from the rate creates a confidence interval indicating that the true rate lies between the lower and upper bounds of this interval with 95 % statistical confidence.

Postneonatal Death: Deaths occurring to individuals 28–364 days of age.
 -: A dash indicates quantity zero.

Source: 2007-2017 Michigan Resident Death Files,
 Division for Vital Records & Health Statistics, Michigan Department of Health & Human Services